


## Glycine-1-<sup>13</sup>C, <sup>15</sup>N

<b>Cat. No.:</b>	HY-Y0966S5
<b>CAS No.:</b>	112898-03-0
<b>Molecular Formula:</b>	C <sup>13</sup> H <sub>5</sub> <sup>15</sup> N <sub>2</sub> O <sub>2</sub>
<b>Molecular Weight:</b>	77.05
<b>Target:</b>	iGluR; Endogenous Metabolite; Isotope-Labeled Compounds
<b>Pathway:</b>	Membrane Transporter/Ion Channel; Neuronal Signaling; Metabolic Enzyme/Protease; Others
<b>Storage:</b>	Please store the product under the recommended conditions in the Certificate of Analysis.



### BIOLOGICAL ACTIVITY

<b>Description</b>	Glycine-1- <sup>13</sup> C, <sup>15</sup> N is the <sup>13</sup> C- and <sup>15</sup> N-labeled Glycine. Glycine is an inhibitory neurotransmitter in the CNS and also acts as a co-agonist along with glutamate, facilitating an excitatory potential at the glutaminergic N-methyl-D-aspartic acid (NMDA) receptors.
<b>IC<sub>50</sub> &amp; Target</b>	NMDA Receptor
<b>In Vitro</b>	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

[1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother*. 2019;53(2):211-216.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA