


Glycine-¹⁵N

Cat. No.:	HY-Y0966S
CAS No.:	7299-33-4
Molecular Formula:	C ₂ H ₃ ¹⁵ N ₂ O ₂
Molecular Weight:	76.06
Target:	Endogenous Metabolite; iGluR
Pathway:	Metabolic Enzyme/Protease; Membrane Transporter/Ion Channel; Neuronal Signaling
Storage:	4°C, protect from light, stored under nitrogen * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light, stored under nitrogen)



BIOLOGICAL ACTIVITY

Description	Glycine- ¹⁵ N is the ¹⁵ N-labeled Glycine. Glycine is an inhibitory neurotransmitter in the CNS and also acts as a co-agonist along with glutamate, facilitating an excitatory potential at the glutamergic N-methyl-D-aspartic acid (NMDA) receptors[1].
IC ₅₀ & Target	NMDA Receptor
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother*. 2019;53(2):211-216.
- [2]. Johnson JW, et al. Glycine potentiates the NMDA response in cultured mouse brain neurons. *Nature*. 1987 Feb 5-11;325(6104):529-31.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA