


PEG1500

Cat. No.:	HY-Y0873E		
CAS No.:	25322-68-3		
Molecular Weight:	1500		
Target:	Biochemical Assay Reagents		
Pathway:	Others		
Storage:	Pure form	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

H₂O : 100 mg/mL (66.67 mM; Need ultrasonic)

Concentration	Solvent	Mass	1 mg	5 mg	10 mg
			1 mM	0.6667 mL	3.3333 mL
5 mM			0.1333 mL	0.6667 mL	1.3333 mL
10 mM			0.0667 mL	0.3333 mL	0.6667 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

PEG1500 can be used as an excipient, such as ointment base, lubricant. Pharmaceutical excipients, or pharmaceutical auxiliaries, refer to other chemical substances used in the pharmaceutical process other than pharmaceutical ingredients. Pharmaceutical excipients generally refer to inactive ingredients in pharmaceutical preparations, which can improve the stability, solubility and processability of pharmaceutical preparations. Pharmaceutical excipients also affect the absorption, distribution, metabolism, and elimination (ADME) processes of co-administered drugs^[1].

REFERENCES

[1]. Yang J, Shen MH. Polyethylene glycol-mediated cell fusion. *Methods Mol Biol.* 2006;325:59-66.

[2]. Elder DP, et al. Pharmaceutical excipients - quality, regulatory and biopharmaceutical considerations. *Eur J Pharm Sci.* 2016 May 25;87:88-99.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA