

## Sodium metabisulfite

<b>Cat. No.:</b>	HY-Y1326
<b>CAS No.:</b>	7681-57-4
<b>Molecular Formula:</b>	Na <sub>2</sub> O <sub>3</sub> S <sub>2</sub>
<b>Molecular Weight:</b>	190.11
<b>Target:</b>	Biochemical Assay Reagents
<b>Pathway:</b>	Others
<b>Storage:</b>	4°C, sealed storage, away from moisture * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



### SOLVENT & SOLUBILITY

#### In Vitro

H<sub>2</sub>O : 100 mg/mL (526.01 mM; Need ultrasonic)  
DMSO : 4 mg/mL (21.04 mM; ultrasonic and warming and heat to 60°C)

Preparing Stock Solutions	Solvent	Mass	1 mg	5 mg	10 mg
	Concentration				
	1 mM		5.2601 mL	26.3006 mL	52.6011 mL
	5 mM		1.0520 mL	5.2601 mL	10.5202 mL
	10 mM		0.5260 mL	2.6301 mL	5.2601 mL

Please refer to the solubility information to select the appropriate solvent.

### BIOLOGICAL ACTIVITY

#### Description

Sodium metabisulfite can be used as an excipient, such as antibacterial agent, preservative, antioxidant. Pharmaceutical excipients, or pharmaceutical auxiliaries, refer to other chemical substances used in the pharmaceutical process other than pharmaceutical ingredients. Pharmaceutical excipients generally refer to inactive ingredients in pharmaceutical preparations, which can improve the stability, solubility and processability of pharmaceutical preparations. Pharmaceutical excipients also affect the absorption, distribution, metabolism, and elimination (ADME) processes of co-administered drugs [1].

### REFERENCES

[1]. Elder DP, et al. Pharmaceutical excipients - quality, regulatory and biopharmaceutical considerations. Eur J Pharm Sci. 2016 May 25;87:88-99.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA