2-(Azido-PEG3-amido)-1,3-bis(NHS ester)

Cat. No.: CAS No.: Molecular Formula: Molecular Weight: Target: Pathway: Storage:	HY-140865 2320560-36-7 C ₂₆ H ₃₈ N ₆ O ₁₄ 658.61 PROTAC Linkers PROTAC Please store the product under the recommended conditions in the Certificate of Analysis.	$N^{N^{N^{N^{N^{N^{N^{N^{N^{N^{N^{N^{N^{N$
---	---	--

BIOLOGICALACTIVITY		
BIOLOGICAL ACTIVITY		
Description	2-(Azido-PEG3-amido)-1,3-bis(NHS ester) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs ^[1] . 2- (Azido-PEG3-amido)-1,3-bis(NHS ester) is a click chemistry reagent, it contains an Azide group and can undergo copper- catalyzed azide-alkyne cycloaddition reaction (CuAAc) with molecules containing Alkyne groups. Strain-promoted alkyne- azide cycloaddition (SPAAC) can also occur with molecules containing DBCO or BCN groups.	
IC ₅₀ & Target	PEGs Alkyl/ether	
In Vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909

609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

