# RedChemExpress

# Product Data Sheet

# Inhibitors • Screening Libraries • Proteins

## (5,6)TAMRA-PEG3-Azide-PEG3-Desthiobiotin

Cat. No.:	HY-140948	
Molecular Formula:	C <sub>57</sub> H <sub>81</sub> N <sub>11</sub> O <sub>19</sub>	Man and a start of the start of
Molecular Weight:	1144.34	o the
Target:	Fluorescent Dye	WWW NO COLONNA HOUSE HIM NO COLONNA HUM HIM O
Pathway:	Others	
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.	what one of the second for the

BIOLOGICAL ACTIVITY		
DIOLOGICAL ACTIV		
Description	(5,6)TAMRA-PEG3-Azide-PEG3-Desthiobiotin is a dye derivative of TAMRA (HY-135640) containing 3 PEG units. (5,6)TAMRA- PEG3-Azide-PEG3-Desthiobiotin contains an Azide group and can undergo copper-catalyzed azide-alkyne cycloaddition reaction (CuAAc) with molecules containing Alkyne groups. Strain-promoted alkyne-azide cycloaddition (SPAAC) can also occur with molecules containing DBCO or BCN groups.	
IC <sub>50</sub> & Target	PEGs	
In Vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

### REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

### Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA