6-Methoxypurine-9-beta-D-(3-methoxy riboside)

Cat. No.: HY-152647

Molecular Formula: $C_{12}H_{16}N_4O_5$

Molecular Weight: 296.28

Target: Nucleoside Antimetabolite/Analog

Pathway: Cell Cycle/DNA Damage

Storage: Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description

6-Methoxypurine-9-beta-D-(3-methoxy riboside) is a purine nucleoside analog. Purine nucleoside analogs have broad antitumor activity targeting indolent lymphoid malignancies. Anticancer mechanisms in this process rely on inhibition of DNA synthesis, induction of apoptosis, etc[1].

REFERENCES

[1]. Virág L, Szabó C. Purines inhibit poly(ADP-ribose) polymerase activation and modulate oxidant-induced cell death. FASEB J. 2001 Jan;15(1):99-107.

[2]. Saugstad OD. Hypoxanthine as an indicator of hypoxia: its role in health and disease through free radical production. Pediatr Res. 1988 Feb;23(2):143-50.

[3]. Robak T, Robak P. Purine nucleoside analogs in the treatment of rarer chronic lymphoid leukemias. Curr Pharm Des. 2012;18(23):3373-88.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA