


## 7BIO

Cat. No.:	HY-121035
CAS No.:	916440-85-2
Molecular Formula:	C <sub>16</sub> H <sub>10</sub> BrN <sub>3</sub> O <sub>2</sub>
Molecular Weight:	356.17
Target:	CDK; GSK-3
Pathway:	Cell Cycle/DNA Damage; PI3K/Akt/mTOR; Stem Cell/Wnt
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



### SOLVENT & SOLUBILITY

In Vitro	DMSO : 250 mg/mL (701.91 mM; Need ultrasonic)						
	Ethanol : 50 mg/mL (140.38 mM; Need ultrasonic)						
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	5 mg	10 mg	
				1 mM	2.8076 mL	14.0382 mL	28.0765 mL
				5 mM	0.5615 mL	2.8076 mL	5.6153 mL
10 mM				0.2808 mL	1.4038 mL	2.8076 mL	
Please refer to the solubility information to select the appropriate solvent.							
In Vivo	1. Add each solvent one by one: 10% EtOH >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (7.02 mM); Clear solution						
	2. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (5.84 mM); Clear solution						

### BIOLOGICAL ACTIVITY

Description	7BIO (7-Bromoindirubin-3-Oxime) is the derivate of indirubin. 7BIO (7-Bromoindirubin-3-Oxime) has inhibitory effects against cyclin-dependent kinase-5 (CDK5) and glycogen synthase kinase-3β (GSK3β). 7BIO (7-Bromoindirubin-3-Oxime) inhibits Aβ oligomer-induced neuroinflammation, synaptic impairments, tau hyper-phosphorylation, activation of astrocytes and microglia, and attenuates Aβ oligomer-induced cognitive impairments in mice <sup>[1]</sup> .	
IC <sub>50</sub> & Target	CDK5	GSK3β
In Vitro	7BIO (1 and 10 μM; 24 hours) has neuroprotective effects and can prevent, but cannot rescue Aβ <sub>1-42</sub> oligomer-induced cell death in SH-SY5Y cells <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

	Cell Viability Assay <sup>[1]</sup>
Cell Line:	SH-SY5Y cells
Concentration:	1 and 10 $\mu$ M
Incubation Time:	24 hours
Result:	Inhibit A $\beta$ oligomer-induced neuronal death. 7BIO prevented, but did not rescue A $\beta$ 1-42 oligomer-induced cell death in SH-SY5Y cells.
<b>In Vivo</b>	<p>7Bio (2.3, 7.0, and 23.3 <math>\mu</math>g/kg; bilateral ventricle injection) significantly attenuates A<math>\beta</math> oligomer-induced impairment of recognition, spatial learning and memory in mice<sup>[1]</sup>.</p> <p>7Bio (2.3, 7.0, and 23.3 <math>\mu</math>g/kg; bilateral ventricle injection) decreases A<math>\beta</math> oligomer-induced increase of TNF-<math>\alpha</math> and IL-6 production in the brain and the expression of synapsin-1 and PSD-95 in the hippocampal region of mice<sup>[1]</sup>.</p> <p>7Bio (2.3, 7.0, and 23.3 <math>\mu</math>g/kg; bilateral ventricle injection) attenuates A<math>\beta</math> oligomer-induced increase expression of pTau, activation of microglia and astrogliosis in the brain of mice<sup>[1]</sup>.</p> <p>7Bio (2.3, 7.0, and 23.3 <math>\mu</math>g/kg; bilateral ventricle injection) prevents decreased expression of pSer9-GSK3<math>\beta</math> and has no significant effects on the Tau protein level<sup>[1]</sup>.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>
Animal Model:	8 weeks mice (30 g) <sup>[1]</sup>
Dosage:	2.3, 7.0, and 23.3 $\mu$ g/kg
Administration:	bilateral ventricle injection
Result:	Attenuates A $\beta$ oligomer-induced impairment of recognition, spatial learning and memory in mice.

## REFERENCES

[1]. Chen L, et al. Indirubin Derivative 7-Bromoindirubin-3-Oxime (7Bio) Attenuates A $\beta$  Oligomer-Induced Cognitive Impairments in Mice. *Front Mol Neurosci.* 2017;10:393.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA