


Amine-PEG-thiol (MW 2000)

Cat. No.:	HY-140651		
Target:	PROTAC Linkers		
Pathway:	PROTAC		
Storage:	Powder	-20°C	3 years
	In solvent	-80°C	6 months
		-20°C	1 month



MW 2000

SOLVENT & SOLUBILITY

In Vitro	DMSO : 50 mg/mL (Need ultrasonic)
In Vivo	<ol style="list-style-type: none"> Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 0.83 mg/mL (Infinity mM); Clear solution Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 0.83 mg/mL (Infinity mM); Clear solution

BIOLOGICAL ACTIVITY

Description	Amine-PEG-thiol (MW 2000) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs ^[1] .
IC₅₀ & Target	PEGs
In Vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA