


Aminoxy-PEG3-C2-thiol

Cat. No.:	HY-134512		
CAS No.:	1200365-63-4		
Molecular Formula:	C ₈ H ₁₉ NO ₄ S		
Molecular Weight:	225.31		
Target:	PROTAC Linkers		
Pathway:	PROTAC		
Storage:	Powder	-20°C	3 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (443.83 mM; Need ultrasonic)					
	Preparing Stock Solutions	Solvent Concentration	Mass			
			1 mg	5 mg	10 mg	
			1 mM	4.4383 mL	22.1916 mL	44.3833 mL
			5 mM	0.8877 mL	4.4383 mL	8.8767 mL
10 mM	0.4438 mL	2.2192 mL	4.4383 mL			
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (11.10 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (11.10 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (11.10 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	Aminoxy-PEG3-C2-thiol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs ^[1] .
IC ₅₀ & Target	PEGs
In Vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA