# Arimoclomol citrate

MedChemExpress

Cat. No.:HY-106443BCAS No.:368860-21-3Molecular Formula:C <sub>20</sub> H <sub>28</sub> ClN <sub>3</sub> O <sub>10</sub> Molecular Weight:505.9Target:HSPPathway:Cell Cycle/DNA Damage; Metabolic Enzyme/ProteaseStorage:Please store the product under the recommended conditions in the Certificate of Analysis.			
Molecular Formula: $C_{20}H_{28}ClN_3O_{10}$ $O \longrightarrow Cl \longrightarrow C$	Cat. No.:	HY-106443B	
Molecular Weight:       505.9         Target:       HSP         Pathway:       Cell Cycle/DNA Damage; Metabolic Enzyme/Protease         Storage:       Please store the product under the recommended conditions in the Certificate of	CAS No.:	368860-21-3	
Target:       HSP         Pathway:       Cell Cycle/DNA Damage; Metabolic Enzyme/Protease         Storage:       Please store the product under the recommended conditions in the Certificate of	Molecular Formula:	C <sub>20</sub> H <sub>28</sub> ClN <sub>3</sub> O <sub>10</sub>	
Pathway:       Cell Cycle/DNA Damage; Metabolic Enzyme/Protease         Storage:       Please store the product under the recommended conditions in the Certificate of	Molecular Weight:	505.9	сі он
Storage: Please store the product under the recommended conditions in the Certificate of	Target:	HSP	o <sup>o</sup> ∽ <sup>oh</sup> o
Storage: Please store the product under the recommended conditions in the Certificate of	Pathway:	Cell Cycle/DNA Damage; Metabolic Enzyme/Protease	но
	Storage:		ОН

BIOLOGICAL ACTIVITY				
Description	Arimoclomol citrate (B neurons by enhancing	Arimoclomol citrate (BRX-220 citrate) is a co-inducer of heat shock proteins (HSP) <sup>[1]</sup> . Arimoclomol citrate protects motor neurons by enhancing Hsp expression, thus directly affecting protein aggregation and clearance of misfolded assemblies via the proteasome-ubiquitin system <sup>[2]</sup> .		
IC <sub>50</sub> & Target	HSP			
In Vivo	(CCK)-induced acute pa	Arimoclomol (BRX-220; 20 mg/kg; orally administered for 5 d) has a protective effect against cholecystokinin-octapeptide (CCK)-induced acute pancreatitis <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.		
	Animal Model:	Male Wistar rats weighing 240 to 270 $g^{[1]}$		
	Dosage:	20 mg/kg		
	Administration:	Administered intragastrically, for 5 days		
	Result:	Had a protective effect against CCK-induced acute pancreatitis.		

## CUSTOMER VALIDATION

- Exp Eye Res. 2019 Nov;188:107792.
- University of Graz. Institute of Pharmaceutical Science. 2021 Nov.

See more customer validations on <u>www.MedChemExpress.com</u>

### REFERENCES

[1]. Rakonczay Z Jr, et al. Nontoxic heat shock protein coinducer BRX-220 protects against acute pancreatitis in rats. Free Radic Biol Med. 2002 Jun 15;32(12):1283-92.

[2]. Phukan J. Arimoclomol, a coinducer of heat shock proteins for the potential treatment of amyotrophic lateral sclerosis. IDrugs. 2010 Jul;13(7):482-96.

#### Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA