


Azido-PEG2-alcohol

Cat. No.:	HY-140797		
CAS No.:	139115-90-5		
Molecular Formula:	C ₄ H ₉ N ₃ O ₂		
Molecular Weight:	131.13		
Target:	PROTAC Linkers		
Pathway:	PROTAC		
Storage:	Pure form	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



SOLVENT & SOLUBILITY

In Vitro

H₂O : 100 mg/mL (762.60 mM; Need ultrasonic)

Concentration	Mass		
	1 mg	5 mg	10 mg
1 mM	7.6260 mL	38.1301 mL	76.2602 mL
5 mM	1.5252 mL	7.6260 mL	15.2520 mL
10 mM	0.7626 mL	3.8130 mL	7.6260 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

Azido-PEG2-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs^[1]. Azido-PEG2-alcohol is a click chemistry reagent, it contains an Azide group and can undergo copper-catalyzed azide-alkyne cycloaddition reaction (CuAAC) with molecules containing Alkyne groups. Strain-promoted alkyne-azide cycloaddition (SPAAC) can also occur with molecules containing DBCO or BCN groups.

IC₅₀ & Target

PEGs

In Vitro

PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA