


Azido-PEG3-NHS ester

Cat. No.:	HY-140764	
CAS No.:	1092654-47-1	
Molecular Formula:	C ₁₂ H ₁₈ N ₄ O ₇	
Molecular Weight:	330.29	
Target:	PROTAC Linkers	
Pathway:	PROTAC	
Storage:	Pure form	-20°C 3 years
	In solvent	-80°C 6 months
		-20°C 1 month



SOLVENT & SOLUBILITY

In Vitro

Ethanol : 50 mg/mL (151.38 mM; Need ultrasonic)

Concentration	Solvent	Mass		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	3.0276 mL	15.1382 mL	30.2764 mL
	5 mM	0.6055 mL	3.0276 mL	6.0553 mL
	10 mM	0.3028 mL	1.5138 mL	3.0276 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

Azido-PEG3-NHS ester is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs^[1]. Azido-PEG3-NHS ester is a click chemistry reagent, it contains an Azide group and can undergo copper-catalyzed azide-alkyne cycloaddition reaction (CuAAC) with molecules containing Alkyne groups. Strain-promoted alkyne-azide cycloaddition (SPAAC) can also occur with molecules containing DBCO or BCN groups.

IC₅₀ & Target

PEGs

In Vitro

PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins^[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA