


## Azido-PEG9-alcohol

<b>Cat. No.:</b>	HY-140800		
<b>CAS No.:</b>	1984776-37-5		
<b>Molecular Formula:</b>	C <sub>18</sub> H <sub>37</sub> N <sub>3</sub> O <sub>9</sub>		
<b>Molecular Weight:</b>	439.5		
<b>Target:</b>	PROTAC Linkers		
<b>Pathway:</b>	PROTAC		
<b>Storage:</b>	Pure form	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### BIOLOGICAL ACTIVITY

<b>Description</b>	Azido-PEG9-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs <sup>[1]</sup> . Azido-PEG9-alcohol is a click chemistry reagent, it contains an Azide group and can undergo copper-catalyzed azide-alkyne cycloaddition reaction (CuAAC) with molecules containing Alkyne groups. Strain-promoted alkyne-azide cycloaddition (SPAAC) can also occur with molecules containing DBCO or BCN groups.
<b>IC<sub>50</sub> &amp; Target</b>	PEGs
<b>In Vitro</b>	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA