Boc-MLF TFA

Cat. No.:	HY-103473A	
Molecular Formula:	$C_{27}H_{40}F_{3}N_{3}O_{8}S$	S
Molecular Weight:	623.68	
Target:	Formyl Peptide Receptor (FPR)	
Pathway:	GPCR/G Protein	
Storage:	4°C, sealed storage, away from moisture * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)	HO F F

SOLVENT & SOLUBILITY

In Vitro	-	DMSO : 150 mg/mL (240.51 mM; Need ultrasonic) H ₂ O : < 0.1 mg/mL (insoluble)					
		Solvent Mass Concentration	1 mg	5 mg	10 mg		
	Preparing Stock Solutions	1 mM	1.6034 mL	8.0169 mL	16.0339 mL		
		5 mM	0.3207 mL	1.6034 mL	3.2068 mL		
		10 mM	0.1603 mL	0.8017 mL	1.6034 mL		
	Please refer to the so	Please refer to the solubility information to select the appropriate solvent.					
n Vivo		1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (4.01 mM); Clear solution					
		2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (4.01 mM); Clear solution					
		3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (4.01 mM); Clear solution					

Boc-MLF (TFA) is a peptide, used as a specific formyl peptide receptor (FPR) antagonist, also inhibits the signaling through

Boc-MLF inhibits superoxide production induced by FPR agonist fMLF, with an EC_{50} of 630 nM in neutrophils^[1].

?Boc-MLF (25 μM) blocks FPRL1-agonist serum amyloid A (SAA) induced calcium response^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

formyl peptide receptor like 1 (FPRL1) at higher concentrations^[1].

Inhibitors

•

Screening Libraries

•

Proteins

Description

IC₅₀ & Target

In Vitro

BIOLOGICAL ACTIVITY

FPR/FPRL1^[1]

CUSTOMER VALIDATION

- World J Gastroenterol. 2023 Jun 14, 29(22): 3422-3439.
- SSRN. 2021 Mar 24.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Stenfeldt AL, et al. Cyclosporin H, Boc-MLF and Boc-FLFLF are antagonists that preferentially inhibit activity triggered through the formyl peptide receptor. Inflammation. 2007 Dec;30(6):224-9. Epub 2007 Aug 9.

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA