Product Data Sheet

DHODH-IN-17

Molecular Weight:

Cat. No.: HY-128068 CAS No.: 16344-26-6

Molecular Formula: $C_{12}H_9CIN_2O_2$

Dihydroorotate Dehydrogenase Target: Pathway: Metabolic Enzyme/Protease

248.67

Storage: Powder -20°C 3 years

> 4°C In solvent -80°C 6 months

> > -20°C 1 month

2 years

SOLVENT & SOLUBILITY

In Vitro

DMSO: 83.33 mg/mL (335.10 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	4.0214 mL	20.1070 mL	40.2139 mL
	5 mM	0.8043 mL	4.0214 mL	8.0428 mL
	10 mM	0.4021 mL	2.0107 mL	4.0214 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- 1. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (8.36 mM); Clear solution
- 2. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 1.67 mg/mL (6.72 mM); Clear solution
- 3. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 1.67 mg/mL (6.72 mM); Clear solution

BIOLOGICAL ACTIVITY

Description	DHODH-IN-17, a 2-anilino nicotinic acid, is a human DHODH inhibitor (IC $_{50}$ =0.40 μ M). DHODH-IN-17 can be used for theresearch of acute myeloid leukemia (AML) ^[1] .			
IC ₅₀ & Target	IC50: 0.4 μM (DHODH) ^[1]			
In Vitro	DHODH-IN-17, a 2-anilino nicotinic acid, is a human DHODH inhibitor with an IC $_{50}$ value of 0.40 μ M $^{[1]}$. MCE has not independently confirmed the accuracy of these methods. They are for reference only.			

REFERENCES						
	ent of ML390: A Human DHOI	DH Inhibitor That Induces Differe	ntiation in Acute Myeloid Leukemia. ACS Me	d Chem Lett. 2016;7(12):1112-1117.		
	Caution: Product has not been fully validated for medical applications. For research use only.					
	Tel: 609-228-6898	Fax: 609-228-5909	E-mail: tech@MedChemExpress.c	com		
	Address:	1 Deer Park Dr, Suite Q, Monm	nouth Junction, NJ 08852, USA			

Page 2 of 2 www.MedChemExpress.com