Proteins

Inhibitors

Screening Libraries

DiBAC4(3)

Cat. No.: HY-101892 CAS No.: 70363-83-6 Molecular Formula: $C_{27}H_{40}N_4O_6$ Molecular Weight: 516.63

Target: Fluorescent Dye

Pathway: Others

4°C, protect from light Storage:

* In solvent: -80°C, 6 months; -20°C, 1 month (protect from light)

Product Data Sheet

SOLVENT & SOLUBILITY

In Vitro

DMSO: 50 mg/mL (96.78 mM; Need ultrasonic)

| Preparing Stock Solutions | Solvent Mass Concentration | 1 mg | 5 mg | 10 mg |
|------------------------------|-------------------------------|-----------|-----------|------------|
| | 1 mM | 1.9356 mL | 9.6781 mL | 19.3562 mL |
| | 5 mM | 0.3871 mL | 1.9356 mL | 3.8712 mL |
| | 10 mM | 0.1936 mL | 0.9678 mL | 1.9356 mL |

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

DiBAC4(3) is a voltage-sensitive fluorescent dye (λ_{ex} =490 nm, λ_{em} =505 nm).

In Vitro

The membrane hyperpolarization induced by 10 μM Evans blue (EB) in HEKBKα is clearly observed with DiBAC4(3), while the change in membrane potential (MP) by addition of 3 mM tetraethylammonium chloride (TEA) appears more slowly than that measured with microelectrode. The time to peak of hyperpolarization is 2.3±0.9 s (n=4) and 35.0±2.6 s (n=12, P<0.01) by the measurements with microelectrodes and DiBAC4(3), respectively^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Cell Assay [1]

Prior to the fluorescence measurements, cells are incubated in KRH (Krebs-Ringer-HEPES) buffer containing with 100 nM DiBAC4(3) for 20 min at room temperature. The stained cells are used for experiments without washing. The fluorescence emission is collected using a 505 nm dicroic mirror and a BA filter (>520 nm)^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Cell Metab. 2022 Sep 7;S1550-4131(22)00359-X.
- ACS Omega. 2023 Mar 6.
- J Cancer Res Clin Oncol. 2022 Nov 27.
- Research Square Preprint. 2020 Nov.

See more customer validations on $\underline{www.\mathsf{MedChemExpress.com}}$

| _ | _ | _ | _ | _ | _ | | - | _ |
|---|---|---|---|---|---|----|---|----|
| D | _ | _ | _ | u | _ | NΙ | | FS |
| | | | | | | | | |

[1]. Yamada A, et al. Usefulness and limitation of DiBAC4(3), a voltage-sensitive fluorescent dye, for the measurement of membrane potentials regulated by recombinant large conductance Ca2+-activated K+ channels in HEK293 cells. Jpn J Pharmacol. 2001 Jul;86(3):342-50.

 $\label{lem:caution:Product} \textbf{Caution: Product has not been fully validated for medical applications. For research use only.}$

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA