## Diclofenac epolamine

MedChemExpress

Cat. No.:	HY-109547	O
CAS No.:	119623-66-4	сі П СІ СІ
Molecular Formula:	C <sub>20</sub> H <sub>24</sub> Cl <sub>2</sub> N <sub>2</sub> O <sub>3</sub>	N N
Molecular Weight:	411.32	
Target:	СОХ	✓ `CI <>
Pathway:	Immunology/Inflammation	ОН
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.	$\langle N \rangle$

BIOLOGICAL AC			
Description	Flector Patch (Diclofen arthritis Pain, acute Pa	Flector Patch (Diclofenac epolamine) is a non-steroidal anti-inflammatory drug (NSAID) <sup>[1]</sup> . Flector Patch is used for relieving arthritis Pain, acute Pain, osteoarthritis, and actinic Keratosis. Flector Patch has good skin absorption characteristics without local adverse reactions and allergies <sup>[2]</sup> .	
In Vitro	Flector Patch (Diclofen and sustains to 12 h af	Flector Patch (Diclofenac epolamine) (1.3% w/w, 24 hours) increases diclofenac partitioning into the skin <sup>[1]</sup> . Flector Patch (Diclofenac epolamine) (1.3% w/w) loaded by Microemulsion (ME) enhances the transport of drug through skin and sustains to 12 h after removal of ME <sup>[2]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	
In Vivo	[2]	Flector Patch (Diclofenac epolamine) (1.3% w/w, exteral application) displays effective in treating inflammatory conditions [2]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.	
	Animal Model:	Rat paw edema inflammation model (induced with carrageenan) <sup>[2]</sup>	
	Dosage:	1.3% w/w	
	Administration:	Exteral application	
	Result:	Inhibited edema and showed anti-inflammatory efficacy at 6 hours after application.	

## REFERENCES

[1]. Del Río-Sancho S, et al. Topical delivery of diclofenac into and across equine skin from a novel liquid diclofenac epolamine formulation. J Vet Pharmacol Ther. 2016 Dec. 39(6):578-583.

[2]. Fouad SA, Basalious EB, El-Nabarawi MA, Tayel SA. Microemulsion and poloxamer microemulsion-based gel for sustained transdermal delivery of diclofenac epolamine using in-skin drug depot: in vitro/in vivo evaluation. Int J Pharm. 2013 Sep 10;453(2):569-78.

## Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA