


## Dihydrorhodamine 123

<b>Cat. No.:</b>	HY-101894
<b>CAS No.:</b>	109244-58-8
<b>Molecular Formula:</b>	C <sub>21</sub> H <sub>18</sub> N <sub>2</sub> O <sub>3</sub>
<b>Molecular Weight:</b>	346.38
<b>Target:</b>	Fluorescent Dye
<b>Pathway:</b>	Others
<b>Storage:</b>	-20°C, protect from light

\* The compound is unstable in solutions, freshly prepared is recommended.



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 100 mg/mL (288.70 mM; Need ultrasonic)				
		Solvent Concentration	Mass		
	<b>Preparing Stock Solutions</b>		1 mg	5 mg	10 mg
		1 mM	2.8870 mL	14.4350 mL	28.8700 mL
		5 mM	0.5774 mL	2.8870 mL	5.7740 mL
	10 mM	0.2887 mL	1.4435 mL	2.8870 mL	
Please refer to the solubility information to select the appropriate solvent.					
<b>In Vivo</b>	1. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: 2.5 mg/mL (7.22 mM); Suspended solution; Need ultrasonic				

### BIOLOGICAL ACTIVITY

<b>Description</b>	Dihydrorhodamine 123 (DHR 123) is a fluorescent probe ( $\lambda_{ex}=488$ nm, $\lambda_{em}=525$ nm) <sup>[1]</sup> .
<b>In Vitro</b>	In the presence of 10 $\mu$ M Dihydrorhodamine 123 (DHR 123) the stimulation of the neutrophil NADPH oxidase by the addition of 50 nM phorbol 12-myristate 13-acetat (PMA) results in an increase in the rate of rhodamine generation. The fluorescent intensity of the cells, in the presence of 10 $\mu$ M Dihydrorhodamine 123, increases with time following the addition of 50 nM PMA. In the presence of 10 $\mu$ M Dihydrorhodamine 123, induced HL60 cells show a sustained increase in fluorescence following the addition of 50 nM PMA <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

### PROTOCOL

<b>Cell Assay</b> <sup>[1]</sup>	The HL60 cells are incubated at $6 \times 10^6$ cells/mL in Krebs-Ringer buffer at 37°C containing 10 $\mu$ M Dihydrorhodamine 123
----------------------------------	--

---

(DHR). The generation of  $O_2^-$  is initiated by the addition of 50 nM phorbol 12-myristate 13-acetate (PMA) and the progress of the generation of rhodamine 123 is monitored in 50- $\mu$ L aliquots ( $3 \times 10^5$  cells) diluted tenfold before analysis. The uninduced HL60 cells are loaded with 5  $\mu$ M carboxy SNARF-1 AM acetate (SNARF-AM) in the  $Na^+$  medium for 10 min at 37°C and washed by centrifugation and resuspension to remove unhydrolysed SNARF ester<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

---

## CUSTOMER VALIDATION

- Part Fibre Toxicol. 2022 Mar 29;19(1):24.
- Free Radic Biol Med. 2023 Mar 3;S0891-5849(23)00100-4.
- Commun Biol. 2023 Mar 11;6(1):259.
- J Mol Cell Cardiol. 2021 Jul 2;S0022-2828(21)00135-8.
- Mol Carcinog. 2023 Sep 26.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Lydia M. Henderson et al. Dihydrorhodamine 123: a fluorescent probe for superoxide generation? Eur.J.Biochem. 217, 973-980.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA