


Enterolactone

Cat. No.:	HY-108692	
CAS No.:	78473-71-9	
Molecular Formula:	C ₁₈ H ₁₈ O ₄	
Molecular Weight:	298.33	
Target:	Apoptosis; Endogenous Metabolite	
Pathway:	Apoptosis; Metabolic Enzyme/Protease	
Storage:	Pure form	-20°C 3 years
	In solvent	-80°C 6 months
		-20°C 1 month



BIOLOGICAL ACTIVITY

Description

Enterolactone is a bioactive phenolic metabolite known as a mammalian lignan derived from dietary lignans. Enterolactone has estrogenic properties and anti-breast cancer activity^[1]. Enterolactone is a radiosensitizer for human breast cancer cell lines through impaired DNA repair and increased apoptosis^[2].

In Vitro

Enterolactone (25-75 μM; 48 hours) arrests the growth of MDA-MB-231 breast cancer cells in the 'S' phase^[1]. Enterolactone (25-75 μM; 15 hours) triggers apoptosis in MDA-MB-231 breast cancer cells via caspase-3 activation^[1]. Enterolactone inhibits TGF-β-induced migration of MDA-MB-231 breast cancer cells. Enterolactone inhibits TGF-β-induced invasion of MDA-MB-231 breast cancer cells through ECM. Enterolactone inhibits the TGF-β-induced EMT program in MDA-MB-231 breast cancer cells. Enterolactone reduces the formation of actin stress fibers by inhibiting the expression of CD44 and MAPK-p38. Enterolactone inhibits the ERK/NF-κB/Snail signaling pathway to revert TGF-β-induced EMT in MDA-MB-231 cells^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Cell Viability Assay^[1]

Cell Line:	MDA-MB-231 cells
Concentration:	25, 50, 75 μM
Incubation Time:	48 hours
Result:	There was a non-significant increase (~24%) in the S phase population following treatment with 25 μM EL, whereas there were significant increases (~34% and ~39%) following treatment with 50 and 75 μM EL, respectively.

REFERENCES

[1]. Bigdeli B, et al. Enterolactone: A novel radiosensitizer for human breast cancer cell lines through impaired DNA repair and increased apoptosis. *Toxicol Appl Pharmacol.* 2016;313:180-194.

[2]. Mali AV, et al. Enterolactone modulates the ERK/NF-κB/Snail signaling pathway in triple-negative breast cancer cell line MDA-MB-231 to revert the TGF-β-induced epithelial-mesenchymal transition. *Cancer Biol Med.* 2018;15(2):137-156.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA