


Foretinib

Cat. No.:	HY-10338		
CAS No.:	849217-64-7		
Molecular Formula:	C ₃₄ H ₃₄ F ₂ N ₄ O ₆		
Molecular Weight:	632.65		
Target:	VEGFR; c-Met/HGFR		
Pathway:	Protein Tyrosine Kinase/RTK		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	1 year
		-20°C	6 months



SOLVENT & SOLUBILITY

In Vitro	DMSO : 75 mg/mL (118.55 mM; Need ultrasonic)					
		Solvent Concentration	Mass	1 mg	5 mg	10 mg
	Preparing Stock Solutions	1 mM		1.5807 mL	7.9033 mL	15.8065 mL
		5 mM		0.3161 mL	1.5807 mL	3.1613 mL
10 mM			0.1581 mL	0.7903 mL	1.5807 mL	
Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (3.95 mM); Clear solution					
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (3.95 mM); Clear solution					
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (3.95 mM); Clear solution					

BIOLOGICAL ACTIVITY

Description	Foretinib is a multi-target tyrosine kinase inhibitor with IC ₅₀ s of 0.4 nM and 0.9 nM for Met and KDR.		
IC ₅₀ & Target	KDR	c-Met	
	0.9 nM (IC ₅₀)	0.4 nM (IC ₅₀)	
In Vitro	Foretinib inhibits HGF receptor family tyrosine kinases with IC ₅₀ values of 0.4 nM for Met and 3 nM for Ron. Foretinib also inhibits KDR, Flt-1, and Flt-4 with IC ₅₀ values of 0.9 nM, 6.8 nM and 2.8 nM, respectively. Foretinib inhibits colony growth of		

B16F10, A549 and HT29 cells with IC₅₀ of 40 nM, 29 nM and 165 nM, respectively^[1]. A recent study indicates Foretinib affects cell growth differently in gastric cancer cell lines MKN-45 and KATO-III. Foretinib inhibits phosphorylation of MET and downstream signaling molecules in MKN-45 cells, while targets GFGR2 in KATO-III cells^[2].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

In Vivo

Foretinib (100 mg/kg, p.o.) results in substantial inhibition of phosphorylation of B16F10 tumor Met and ligand (e.g., HGF or VEGF)-induced receptor phosphorylation of Met in liver and Flk-1/KDR in lung, which both persist through 24 hours. Foretinib (30-100 mg/kg, once daily, p.o.) results in reduction in tumor burden. The lung surface tumor burden is reduced by 50% and 58% following treatment with 30 and 100 mg/kg Foretinib, respectively. Foretinib treatment of mice bearing B16F10 solid tumors also results in dose-dependent tumor growth inhibition of 64% and 87% at 30 and 100 mg/kg, respectively. For both studies, administration of Foretinib is well tolerated with no significant body weight loss^[1]. Foretinib is developed to target abnormal signaling of HGF through Met and simultaneously target several receptors tyrosine kinase involved in tumor angiogenesis. Foretinib causes tumor hemorrhage and necrosis in human xenografts within 2 to 4 hours, and maximal tumor necrosis is observed at 96 hours (after five daily doses), resulting in complete regression^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

PROTOCOL

Kinase Assay ^[1]

Kinase inhibition is investigated using one of three assay formats: [³³P]phosphoryl transfer, luciferase-coupled chemiluminescence, or AlphaScreen tyrosine kinase technology. IC₅₀s are calculated by nonlinear regression analysis using XLFit. ³³P-Phosphoryl Transfer Kinase Assay Reactions are performed in 384-well white, clear bottom, high-binding microtiter plates. Plates are coated with 2 μg/well of protein or peptide substrate in a 50 μL volume of coating buffer contained 40 μg/mL substrate (poly(Glu, Tyr) 4:1, 22.5 mM Na₂CO₃, 27.5 mM NaHCO₃, 50 mM NaCl and 3 mM NaN₃). Coated plates are washed once with 50 μL of assay buffer following overnight incubation at room temperature (RT). Test compounds and enzymes are combined with ³³P-γ-ATP (3.3 μCi/nmol) in a total volume of 20 μL. The reaction mixture is incubated at RT for 2 hours and terminated by aspiration. The microtiter plates are subsequently washed 6 times with 0.05% Tween-PBS buffer (PBST). Scintillation fluid (50 μL/well) is added and incorporated ³³P is measured by liquid scintillation spectrometry using a MicroBeta scintillation counter. Luciferase-Coupled Chemiluminescence Assay Reactions are conducted in 384-well white, medium binding microtiter plates. In a first step enzyme and compound are combined and incubated for 60 minutes; reactions are initiated by addition of ATP and peptide substrate (poly(Glu, Tyr) 4:1) in a final volume of 20 μL, and incubated at RT for 2-4 hours. Following the kinase reaction, a 20 μL aliquot of Kinase Glo is added and luminescence signal is measured using a Victor plate reader. Total ATP consumption is limited to 50%. AlphaScreen™ Tyrosine Kinase Assay Donor beads coated with streptavidin and acceptor beads coated with PY100 anti-phosphotyrosine antibody are used. Biotinylated poly(Glu, Tyr) 4:1 is used as the substrate. Substrate phosphorylation is measured by addition of donor/acceptor beads by luminescence following donor-acceptor bead complex formation. Kinase and test compounds are combined and preincubated for 60 minutes, followed by addition of ATP, and biotinylated poly(Glu, Tyr) in a total volume of 20 μL in 384-well white, medium binding microtiter plates. Reaction mixtures are incubated for 1 hour at room temperature. Reactions are quenched by addition of 10 μL of 15-30 μg/mL AlphaScreen bead suspension containing 75 mM Hepes, pH 7.4, 300 mM NaCl, 120 mM EDTA, 0.3% BSA and 0.03% Tween-20. After 2-16 hours incubation at room temperature plates are read using an AlphaQuest reader.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Cell Assay ^[1]

B16F10, A549, and HT29 cells (1.2×10³ per well) are mixed with soft agar and seeded in a 96-well plate containing 10% FBS and EXEL-2880 over a base agar layer. For normoxic conditions, the plates are incubated (37°C) for 12 to 14 days in 21% oxygen, 5% CO₂, and 74% nitrogen, whereas incubation (37°C) under hypoxic conditions is done in a hypoxia chamber in 1% oxygen, 5% CO₂, and 94% nitrogen. The number of colonies is evaluated under each condition following addition of 50% Alamar Blue and fluorescence detection.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Administration ^[1]

In vivo target modulation studies are done in naive mice or mice bearing B16F10 tumors. Foretinib or vehicle (0.9% normal saline) is administered at 10 mL/kg via oral gavage. For examination of Met phosphorylation in liver, HGF (10 μg/mouse) is administered i.v. 10 min before harvest. For examination of Flk-1/KDR phosphorylation in lung, VEGF (10 μg/mouse) is

administered i.v. 30 min before harvest 0.5 h later. Receptor phosphorylation analysis is determined by immunoblot analysis.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Cancer Discov. 2021 Jan;11(1):126-141.
- Nat Biomed Eng. 2018 Aug;2(8):578-588.
- Sci Transl Med. 2018 Jul 18;10(450):eaaq1093.
- Cell Chem Biol. 2018 Feb 15;25(2):206-214.e11.
- RSC Adv. 2019, 9, 4862-4869

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Qian F, et al. Inhibition of tumor cell growth, invasion, and metastasis by EXEL-2880 (XL880, GSK1363089), a novel inhibitor of HGF and VEGF receptor tyrosine kinases. *Cancer Res*, 2009, 69(20), 8009-8016.
- [2]. Kataoka Y, et al. Foretinib (GSK1363089), a multi-kinase inhibitor of MET and VEGFRs, inhibits growth of gastric cancer cell lines by blocking inter-receptor tyrosine kinase networks. *Invest New Drugs*, 2011.
- [3]. Eder JP, et al. A phase I study of foretinib, a multi-targeted inhibitor of c-Met and vascular endothelial growth factor receptor 2. *Clin Cancer Res*, 2010, 16(13), 3507-3516.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA