


Gossypol

Cat. No.:	HY-13407
CAS No.:	303-45-7
Molecular Formula:	C ₃₀ H ₃₀ O ₈
Molecular Weight:	519
Target:	Bcl-2 Family
Pathway:	Apoptosis
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 33.33 mg/mL (64.22 mM; Need ultrasonic)				
	Preparing Stock Solutions	Solvent Concentration	Mass	1 mg	
				5 mg	
				10 mg	
				10 mM	
			1 mg	5 mg	10 mg
	1 mM		1.9268 mL	9.6339 mL	19.2678 mL
	5 mM		0.3854 mL	1.9268 mL	3.8536 mL
	10 mM		0.1927 mL	0.9634 mL	1.9268 mL
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (4.01 mM); Clear solution				

BIOLOGICAL ACTIVITY

Description	Gossypol binds to Bcl-xL protein and Bcl-2 protein with K _i s of 0.5-0.6 μM and 0.2-0.3 mM, respectively.	
IC ₅₀ & Target	Bcl-xL 0.5-0.6 μM (K _i)	Bcl-2 0.2-0.3 mM (K _i)
In Vitro	<p>Gossypol, a natural product isolated from cottonseeds and roots that has been studied as an anticancer agent. The racemic form of Gossypol [(±)-Gossypol] is tested in several clinical trials and is well tolerated. The racemic form Gossypol ((±)-Gossypol) binds to Bcl-xL protein with a K_i of 0.5 to 0.6 μM. (±)-Gossypol also potently binds to Bcl-2 protein with a K_i value of 0.2-0.3 mM. The natural racemic Gossypol has two enantiomers, namely the (-)-Gossypol and (+)-Gossypol enantiomers. The racemic form and each of the enantiomers of Gossypol are tested against UM-SCC-6 and UM-SCC-14A in 6-day MTT assays. (-)-Gossypol exhibits greater growth inhibition relative to (±)-Gossypol than (+)-Gossypol in both cell lines tested (P<0.001). An intermediate growth inhibitory effect is observed with (±)-Gossypol but this effect is only observed at the higher dose of Gossypol (10 μM, P<0.0001)^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>	

PROTOCOL

Cell Assay ^[1]

Two representative UM-SCC cell lines, UM-SCC-6 and UM-SCC-14A, are continuously exposed to 0 (vehicle control), 5 or 10 μ M (\pm)-Gossypol, (-)-Gossypol or (+)-Gossypol in a 6-day MTT cell survival assay^[1].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Clin Transl Med. 2021 Jun;11(6):e467.
- Patent. US20220162561A1.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Oliver CL, et al. In vitro effects of the BH3 mimetic, (-)-Gossypol, on head and neck squamous cell carcinoma cells. Clin Cancer Res. 2004 Nov 15;10(22):7757-63.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA