


Hydroxypyruvic acid lithium hydrate

Cat. No.:	HY-113013A
CAS No.:	209728-15-4
Molecular Formula:	C ₃ H ₃ LiO ₅
Molecular Weight:	128.01
Target:	Endogenous Metabolite
Pathway:	Metabolic Enzyme/Protease
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Hydroxypyruvic acid lithium hydrate (β-Hydroxypyruvic acid lithium hydrate) is an intermediate in the metabolism of glycine, serine and threonine. Hydroxypyruvic acid lithium hydrate is a substrate for serine-pyruvate aminotransferase and glyoxylate reductase/hydroxypyruvate reductase. Hydroxypyruvic acid lithium hydrate is involved in the metabolic disorder which is the dimethylglycine dehydrogenase deficiency pathway.
IC₅₀ & Target	Human Endogenous Metabolite
In Vivo	Hydroxypyruvic acid (lithium hydrate) (intravenous injection; 100 mg/ml; slowly over 10 min) increases the 5-h urinary oxalate and glycolate excretion to 0.68% (6.56 μmol) and 0.53% (5.10 μmol) in control rats, in addition, it increases to 2.43% (23.36 μmol) and 0.79% (7.59 μmol) of the dose in the vitamin-B6-deficient rats ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Teerajetgul Y, et al. Oxalate synthesis from hydroxypyruvate in vitamin-B6-deficient rats. Urol Res. 2007 Aug;35(4):173-8. Epub 2007 Jun 13.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA