


LRRK2 inhibitor 1

Cat. No.:	HY-111493		
CAS No.:	1802525-61-6		
Molecular Formula:	C ₂₀ H ₂₃ N ₅ O ₄		
Molecular Weight:	397.43		
Target:	LRRK2		
Pathway:	Autophagy		
Storage:	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



SOLVENT & SOLUBILITY

In Vitro	DMSO : 62.5 mg/mL (157.26 mM; Need ultrasonic)			
		Solvent Concentration	Mass	
			1 mg	5 mg
	Preparing Stock Solutions		10 mg	
	1 mM	2.5162 mL	12.5808 mL	25.1617 mL
	5 mM	0.5032 mL	2.5162 mL	5.0323 mL
	10 mM	0.2516 mL	1.2581 mL	2.5162 mL
Please refer to the solubility information to select the appropriate solvent.				
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (5.23 mM); Clear solution			
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: 2.08 mg/mL (5.23 mM); Suspended solution; Need ultrasonic			
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (5.23 mM); Clear solution			

BIOLOGICAL ACTIVITY

Description	LRRK2 inhibitor 1 is a potent, selective and oral LRRK2 inhibitor with an pIC ₅₀ of 6.8.
IC₅₀ & Target	pIC ₅₀ : 6.8 nM (LRRK2) ^[1]

REFERENCES

[1]. Ding X, et al. Discovery of 4-ethoxy-7H-pyrrolo[2,3-d]pyrimidin-2-amines as potent, selective and orally bioavailable LRRK2 inhibitors. *Bioorg Med Chem Lett*. 2018 May 15;28(9):1615-1620.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA