


## Lificiguat

<b>Cat. No.:</b>	HY-14927		
<b>CAS No.:</b>	170632-47-0		
<b>Molecular Formula:</b>	C <sub>19</sub> H <sub>16</sub> N <sub>2</sub> O <sub>2</sub>		
<b>Molecular Weight:</b>	304.34		
<b>Target:</b>	Guanylate Cyclase		
<b>Pathway:</b>	GPCR/G Protein		
<b>Storage:</b>	Powder	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	2 years
		-20°C	1 year



### SOLVENT & SOLUBILITY

#### In Vitro

DMSO : ≥ 100 mg/mL (328.58 mM)  
 \* "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	3.2858 mL	16.4290 mL	32.8580 mL
	5 mM	0.6572 mL	3.2858 mL	6.5716 mL
	10 mM	0.3286 mL	1.6429 mL	3.2858 mL

Please refer to the solubility information to select the appropriate solvent.

#### In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline  
 Solubility: ≥ 2.5 mg/mL (8.21 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)  
 Solubility: 2.5 mg/mL (8.21 mM); Suspended solution; Need ultrasonic
- Add each solvent one by one: 10% DMSO >> 90% corn oil  
 Solubility: ≥ 2.5 mg/mL (8.21 mM); Clear solution

### BIOLOGICAL ACTIVITY

#### Description

Lificiguat binds to the β subunit of soluble guanylyl cyclase(sGC) with K<sub>d</sub> of 0.6-1.1 μM in the presence of CO.

#### IC<sub>50</sub> & Target

Kd: 0.6-1.1 μM (sGC, in the presence of CO)<sup>[1]</sup>

#### In Vitro

Soluble guanylate cyclase (sGC) is a heterodimeric heme protein and the primary NO receptor. Lificiguat (YC-1) binds near or directly to the heme-containing domain of the beta subunit. In the absence of CO, Lificiguat (YC-1) binds with K<sub>d</sub>=9-21 μM,

depending on construct. In the presence of CO, these values decrease to 0.6-1.1  $\mu\text{M}$ . Lifigiquat (YC-1) greatly enhanced CO binding to heterodimeric sGC, as expected ( $K_d=1 \mu\text{M}$ ). Lifigiquat (YC-1) stimulates sGC two- to four-fold in the absence of NO but acts synergistically with CO or NO to achieve several hundred fold activation. Binding of Lifigiquat(YC-1) can also overcome inhibitory phosphorylation of sGC<sup>[1]</sup>. Lifigiquat (YC-1) is a soluble guanylyl cyclase (sGC) activator. HCC cell lines HepG2, BEL-7402 and HCCLM3 are incubated for 72 h with Sorafenib and/or Lifigiquat (YC-1). Sorafenib or Lifigiquat (YC-1) alone inhibits HCC cell proliferation in a dose-dependent manner. Moreover, combination of Sorafenib and Lifigiquat (YC-1) significantly suppresses proliferation of HCC cells in a dose-dependent manner. In addition, at the ED<sub>50</sub> doses for both Sorafenib and Lifigiquat (YC-1), combination index values (CI)=0.93 in HepG2, 0.95 in BEL-7402 and 0.72 in HCCLM3 respectively, suggesting that Sorafenib and Lifigiquat (YC-1) synergistically inhibit proliferation of HCC cells<sup>[2]</sup>. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### In Vivo

Lifigiquat (YC-1) (30 or 60 mg/kg, i.p.) inhibits MDA-MB-468 tumor growth in a dose-dependent manner. The effect of the prodrug formulation of Lifigiquat (YC-1), YC-1-S, in MDA-MB-468 tumor-bearing mice is also investigated. In vivo pharmacokinetic analysis reveal that YC-1-S is quickly converted into its active form. Mice are administered 20, 40 or 80 mg/kg YC-1-S p.o. YC-1-S also displays dose-dependent inhibition of MDA-MB468 tumor growth. Both Lifigiquat (YC-1) and YC-1-S dose-dependently reduce tumor weight. Moreover, the mean body weight of mice is not affected by Lifigiquat (YC-1) or YC-1-S compare with vehicle-treated groups<sup>[3]</sup>. Lifigiquat (YC-1) is a potent NO-GC activator reported to improve rodent learning behavior when examined with the Morris water maze (MWM) and avoidance tests. Lifigiquat (YC-1) enhances long-term potentiation (LTP) in hippocampal Schafer collateral-CA1 synapse via the NO-cGMP-PKG-dependent pathway and potentiated LTP induction in the amygdala, increases the activation of ERK, and potentiated the expression of brain-derived neurotrophic factor (BDNF) cAMP response element-binding protein (CREB) in response to fear memory test<sup>[4]</sup>. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

## PROTOCOL

#### Kinase Assay <sup>[1]</sup>

CO dissociation constants are measured by titrating CO from a saturated solution into sGC protein and monitoring the appearance of the CO-bound Soret absorption band. The Ms sGC  $\beta_1$ (1-380) and Bt sGC  $\beta_1$ (1-197) samples are prepared in Ar-purged buffer supplemented with excess dithionite. CO binding experiments are performed in a 10 cm pathlength cuvette for Ms sGC- $\beta_1$ (1-380) and Ms sGC-NT21 using a Cary 50 spectrophotometer with a modified sample holder. Binding data in the presence and absence of 50  $\mu\text{M}$  Lifigiquat (YC-1) is plotted using a single site saturation ligand binding model in SigmaPlot<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Cell Assay <sup>[2]</sup>

Cell proliferation assay is measured using a Cell Counting Kit-8 (CCK-8). Briefly, cells are cultured in 96-well plates at a concentration of  $3 \times 10^3$ /well, incubated for 24 h, and treated with Sorafenib and/or Lifigiquat (YC-1). After 72 h treatment, CCK-8 reagent is added to each well. The absorbance is measured at 450 nm after 2.5 h incubation at 37°C using an automated ELISA plate reader. Any synergistic effects resulting from combination of the compounds are measured using Microsoft Excel software to determine the combination index values (CI>1: antagonistic effect, CI=1: additive effect, and CI<1: synergistic effect)<sup>[2]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Animal Administration <sup>[3][4]</sup>

Mice<sup>[3]</sup>

Fifty-eight female nu/nu mice (4 weeks-old) are used. MDA-MB-468 breast cancer cells ( $5 \times 10^6$  cells per mouse) are suspended in 0.1 mL of Matrigel solution (50% v/v Matrigel in PBS) and inoculated into the mammary fat pads of nude mice. When the tumor masses reach 100 mm<sup>3</sup>, the tumor-bearing mice are randomly divided into groups for treatments with different Lifigiquat (YC-1)/YC-1-S doses. The mice are i.p. injected with YC-1 (30 or 60 mg/kg) or administered YC-1-S p.o. Tumor size and mouse body weight are measured once every 3 days, and tumor volume (mm<sup>3</sup>) is calculated using the equation: length $\times$ (width)<sup>2</sup> $\times$ 0.5. At the end of the experiments, mice are killed and tumor nodules are dissected and weighed. Tumor tissues are subjected to Western blotting.

Rats<sup>[4]</sup>

4-month-old (200-250 g) and 24-month-old (550-600 g) male Wistar-albino rats are used. Lifigiquat (YC-1) is prepared immediately prior to use and given intraperitoneally (i.p.) in a volume of 0.1 mL per 100 g body weight. All rats receives 1

---

mg/kg/day of Lificiguat (YC-1) for 2 weeks. DMSO is administered to 4-month-old and 24-month-old rats (n=10, for each group). Doses are selected to confirm the selected doses on locomotor activity; all results are measured. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

---

## CUSTOMER VALIDATION

- Signal Transduct Target Ther. 2021 Oct 13;6(1):352.
- Adv Healthc Mater. 2022 Dec 5;e2202210.
- Cell Death Dis. 2021 Jan 12;12(1):77.
- Stem Cell Res Ther. 2022 Jul 16;13(1):316.
- Biomed Pharmacother. 2018 Nov;107:1736-1743.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

- [1]. Purohit R, et al. YC-1 binding to the  $\beta$  subunit of soluble guanylyl cyclase overcomes allosteric inhibition by the  $\alpha$  subunit. *Biochemistry*. 2014 Jan 14;53(1):101-14.
- [2]. Kong J, et al. YC-1 enhances the anti-tumor activity of sorafenib through inhibition of signal transducer and activator of transcription 3 (STAT3) in hepatocellular carcinoma. *Mol Cancer*. 2014 Jan 13;13:7.
- [3]. Chang LC, et al. YC-1 inhibits proliferation of breast cancer cells by down-regulating EZH2 expression via activation of c-Cbl and ERK. *Br J Pharmacol*. 2014 Sep;171(17):4010-25.
- [4]. Komsuoglu Celikyurt I, et al. Effects of YC-1 on Learning and Memory Functions of Aged Rats. *Med Sci Monit Basic Res*. 2014 Aug 21;20:130-7.
- 

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA