


Linezolid

Cat. No.:	HY-10394
CAS No.:	165800-03-3
Molecular Formula:	C ₁₆ H ₂₀ FN ₃ O ₄
Molecular Weight:	337
Target:	Bacterial; Antibiotic
Pathway:	Anti-infection
Storage:	4°C, protect from light * In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)



SOLVENT & SOLUBILITY

In Vitro

DMSO : ≥ 100 mg/mL (296.74 mM)
* "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mg	5 mg	10 mg
	1 mM	2.9674 mL	14.8368 mL	29.6736 mL
	5 mM	0.5935 mL	2.9674 mL	5.9347 mL
	10 mM	0.2967 mL	1.4837 mL	2.9674 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 2.5 mg/mL (7.42 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)
Solubility: ≥ 2.5 mg/mL (7.42 mM); Clear solution
- Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 2.5 mg/mL (7.42 mM); Clear solution

BIOLOGICAL ACTIVITY

Description

Linezolid (PNU-100766) is the first member of the class of oxazolidinone synthetic antibiotic. Linezolid acts by inhibiting the initiation of bacterial protein synthesis. Linezolid is used for the treatment of serious infections caused by Gram-positive bacteria that are resistant to several other antibiotics^{[1][2][3]}.

IC₅₀ & Target

Oxazolidinone

In Vitro

Linezolid (PNU-100766) prevents the synthesis of bacterial protein via binding to rRNA on both the 30S and 50S ribosomal subunits^[3].

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

CUSTOMER VALIDATION

- Acta Pharm Sin B. 2017, 52(6): 971-976.
- Int J Biol Macromol. 2023 May 7;124750.
- iScience. 2023 Mar.
- iScience. 5 January 2022, 103731.
- Microb Biotechnol. 2021 Mar 15.

See more customer validations on www.MedChemExpress.com

REFERENCES

- [1]. Clemett D, Markham A. Linezolid. Drugs. 2000 Apr;59(4):815-27; discussion 828.
 - [2]. Chiappini E, Conti C, Galli L et al. Clinical efficacy and tolerability of linezolid in pediatric patients: a systematic review. Clin Ther. 2010 Jan;32(1):66-88.
 - [3]. Perry CM, Jarvis B. Linezolid: a review of its use in the management of serious gram-positive infections. Drugs. 2001;61(4):525-51.
 - [4]. He MZ, Jiang YW, Cai C. Mechanisms and epidemiology of linezolid resistance in staphylococci. Zhonghua Jie He He Hu Xi Za Zhi. 2012 May;35(5):360-2.
 - [5]. Karau MJ, Tilahun AY, Schmidt SM, Clark CR, Patel R, Rajagopalan G. Linezolid is Superior to Vancomycin in Experimental Pneumonia Caused by Superantigen-Producing Staphylococcus aureus in HLA class II Transgenic Mice. Antimicrob Agents Chemother. 2012 Jul 30.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA