


Muscarine chloride

Cat. No.:	HY-121404A
CAS No.:	2303-35-7
Molecular Formula:	C ₉ H ₂₀ ClNO ₂
Molecular Weight:	209.71
Target:	mAChR
Pathway:	GPCR/G Protein; Neuronal Signaling
Storage:	4°C, sealed storage, away from moisture * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



BIOLOGICAL ACTIVITY

Description	Muscarine ((+)-Muscarine) chloride is an agonist of prototype mAChR. Muscarine chloride is a toxin that can stimulate the parasympathetic nervous system ^{[1][2]} .
In Vitro	Muscarine chloride (100 μM) induces an intracellular calcium signal amplitude similar to the one triggered by 10 μM acetylcholine in brain micro vascular endothelial cells (BMVECs) and brain-derived Endothelial cells.3 (bEnd.3) ^[1] . Muscarine chloride (1-30 μM; 2 min) produces a dose-dependent hyperpolarization in a sub-population of the nucleus raphe magnus (NRM) cells that contain 5-hydroxytryptamine (5-HT) on the NRM neurons with an EC ₅₀ value of 2.7 μM ^[2] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Beatrice Mihaela Radu, et al. All muscarinic acetylcholine receptors (M 1-M 5) are expressed in murine brain microvascular endothelium. *Sci Rep.* 2017 Jul 11;7(1):5083.
- [2]. Z Z Pan, et al. Muscarine hyperpolarizes a subpopulation of neurons by activating an M2 muscarinic receptor in rat nucleus raphe magnus in vitro. *J Neurosci.* 1994 Mar;14(3 Pt 1):1332-8.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA