


N-Boc-PEG16-alcohol

| | |
|---------------------------|---|
| Cat. No.: | HY-141194 |
| Molecular Formula: | C ₃₇ H ₇₅ NO ₁₈ |
| Molecular Weight: | 821.99 |
| Target: | PROTAC Linkers |
| Pathway: | PROTAC |
| Storage: | Please store the product under the recommended conditions in the Certificate of Analysis. |



BIOLOGICAL ACTIVITY

| | | |
|-------------------------------------|--|-------------|
| Description | N-Boc-PEG16-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs ^[1] . | |
| IC₅₀ & Target | PEGs | Alkyl/ether |
| In Vitro | PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only. | |

REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA