


## N-Boc-PEG5-bromide

<b>Cat. No.:</b>	HY-120702		
<b>CAS No.:</b>	1392499-32-9		
<b>Molecular Formula:</b>	C <sub>15</sub> H <sub>30</sub> BrNO <sub>6</sub>		
<b>Molecular Weight:</b>	400.31		
<b>Target:</b>	ADC Linker; PROTAC Linkers		
<b>Pathway:</b>	Antibody-drug Conjugate/ADC Related; PROTAC		
<b>Storage:</b>	Pure form	-20°C	3 years
		4°C	2 years
	In solvent	-80°C	6 months
		-20°C	1 month



### BIOLOGICAL ACTIVITY

<b>Description</b>	N-Boc-PEG5-bromide is a PEG/Alkyl/ether-based PROTAC linker can be used in the synthesis of PROTACs. N-Boc-PEG5-bromide is a cleavable ADC linker used in the synthesis of antibody-drug conjugates (ADCs) <sup>[1]</sup> .		
<b>IC<sub>50</sub> &amp; Target</b>	Cleavable Linker	PEGs	Alkyl/ether
<b>In Vitro</b>	<p>PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins. ADCs are comprised of an antibody to which is attached an ADC cytotoxin through an ADC linker.</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p>		

### REFERENCES

[1]. Carston R. Wagner, et al. Protein nanorings. US8236925B1.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA