NH-bis(PEG2-C2-acid)

Cat. No.: CAS No.: Molecular Formula: Molecular Weight: Target: Pathway: Storage:	HY-140514 1919044-99-7 C ₁₄ H ₂₇ NO ₈ 337.37 PROTAC Linkers PROTAC Please store the product under the recommended conditions in the Certificate of Analysis.	но ^щ о о о пределатори
---	--	-----------------------------------

BIOLOGICAL ACTIVITY		
Description	NH-bis(PEG2-C2-acid) is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs ^[1] .	
IC ₅₀ & Target	PEGs	
In Vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562

Caution: Product has not been fully validated for medical applications. For research use only.

Fax: 609-228-5909 E-mail: tech@MedChemExpress.com Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Product Data Sheet



Tel: 609-228-6898