


NLRP3-IN-15

Cat. No.:	HY-149122
CAS No.:	2767369-71-9
Molecular Formula:	C ₂₂ H ₁₉ NO ₄
Molecular Weight:	361.39
Target:	NOD-like Receptor (NLR)
Pathway:	Immunology/Inflammation
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	NLRP3-IN-15 is a potent and selective NLRP3 inflammasome inhibitor. NLRP3-IN-15 inhibits IL-1 β release with an IC ₅₀ of 0.114 μ M. NLRP3-IN-15 can be used for the research of inflammation ^[1] .									
IC₅₀ & Target	NLRP3 inflammasome									
In Vitro	<p>NLRP3-IN-15 (Compound 12a) inhibits IL-1β release with an IC₅₀ of 0.114 μM (ELISA assay)^[1].</p> <p>NLRP3-IN-15 (2 μM, 3 h) inhibits the secretion of IL-1β (p17) and caspase-1 (p20) in mice peritoneal macrophages (PMs)^[1].</p> <p>NLRP3-IN-15 inhibits the formation of the NLRP3 inflammasome complex by inhibiting ASC oligomerization^[1].</p> <p>NLRP3-IN-15 shows metabolic stability in both human and mouse liver microsomes (T_{1/2}: 693 min; Cl_{int}: 2.0 μL/min/mg)^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <p>Western Blot Analysis^[1]</p> <table border="1"> <tr> <td>Cell Line:</td> <td>Mice peritoneal macrophages(PMs)</td> </tr> <tr> <td>Concentration:</td> <td>2 μM</td> </tr> <tr> <td>Incubation Time:</td> <td>3 h</td> </tr> <tr> <td>Result:</td> <td>Inhibited the secretion of IL-1β (p17) and caspase-1 (p20), without affecting pro-IL-1β and pro-caspase-1.</td> </tr> </table>		Cell Line:	Mice peritoneal macrophages(PMs)	Concentration:	2 μ M	Incubation Time:	3 h	Result:	Inhibited the secretion of IL-1 β (p17) and caspase-1 (p20), without affecting pro-IL-1 β and pro-caspase-1.
Cell Line:	Mice peritoneal macrophages(PMs)									
Concentration:	2 μ M									
Incubation Time:	3 h									
Result:	Inhibited the secretion of IL-1 β (p17) and caspase-1 (p20), without affecting pro-IL-1 β and pro-caspase-1.									
In Vivo	<p>NLRP3-IN-15 (Compound 12a) (50 mg/kg; i.p.) shows anti-inflammatory effect in septic mouse model^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <table border="1"> <tr> <td>Animal Model:</td> <td>LPS-induced inflammatory septic mouse model^[1]</td> </tr> <tr> <td>Dosage:</td> <td>50 mg/kg</td> </tr> <tr> <td>Administration:</td> <td>i.p.</td> </tr> <tr> <td>Result:</td> <td>Decreased the release of IL-1β in mouse serum. Relieved thickening of the alveolar wall in lung.</td> </tr> </table>		Animal Model:	LPS-induced inflammatory septic mouse model ^[1]	Dosage:	50 mg/kg	Administration:	i.p.	Result:	Decreased the release of IL-1 β in mouse serum. Relieved thickening of the alveolar wall in lung.
Animal Model:	LPS-induced inflammatory septic mouse model ^[1]									
Dosage:	50 mg/kg									
Administration:	i.p.									
Result:	Decreased the release of IL-1 β in mouse serum. Relieved thickening of the alveolar wall in lung.									

Animal Model:	Mice ^[1]
Dosage:	20 mg/kg
Administration:	p.o. or i.p.
Result:	Pharmacokinetic profile of NLRP3-IN-15 (compound 12a).

parameter	dose (mg/kg)	T _{1/2} (h)	T _{max} (h)	F (%)
PO	20	1.659	0.5	9.6
IP	20	1.807	0.167	21.2

REFERENCES

[1]. Li J, et al. Scaffold Hybrid of the Natural Product Tanshinone I with Piperidine for the Discovery of a Potent NLRP3 Inflammasome Inhibitor. J Med Chem. 2023 Feb 23;66(4):2946-2963.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA