


## Nicotinamide riboside chloride

<b>Cat. No.:</b>	HY-123033A
<b>CAS No.:</b>	23111-00-4
<b>Molecular Formula:</b>	C <sub>11</sub> H <sub>15</sub> ClN <sub>2</sub> O <sub>5</sub>
<b>Molecular Weight:</b>	290.7
<b>Target:</b>	Sirtuin; Endogenous Metabolite
<b>Pathway:</b>	Cell Cycle/DNA Damage; Epigenetics; Metabolic Enzyme/Protease
<b>Storage:</b>	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	H <sub>2</sub> O : 125 mg/mL (430.00 mM; Need ultrasonic)			
	DMSO : 50 mg/mL (172.00 mM; Need ultrasonic)			
		Solvent Concentration	Mass	
	<b>Preparing Stock Solutions</b>		1 mg	5 mg
	1 mM	3.4400 mL	17.1999 mL	34.3997 mL
	5 mM	0.6880 mL	3.4400 mL	6.8799 mL
	10 mM	0.3440 mL	1.7200 mL	3.4400 mL
Please refer to the solubility information to select the appropriate solvent.				
<b>In Vivo</b>	1. Add each solvent one by one: PBS Solubility: 25 mg/mL (86.00 mM); Clear solution; Need ultrasonic			

### BIOLOGICAL ACTIVITY

<b>Description</b>	Nicotinamide riboside Chloride, an orally active NAD <sup>+</sup> precursor, increases NAD <sup>+</sup> levels and activates SIRT1 and SIRT3. Nicotinamide riboside Chloride is a source of vitamin B3 (niacin) and enhances oxidative metabolism, protection against high fat diet-induced metabolic abnormalities <sup>[1]</sup> . Nicotinamide riboside Chloride reduces cognitive deterioration in a transgenic mouse model of Alzheimer's disease <sup>[2]</sup> .		
<b>IC<sub>50</sub> &amp; Target</b>	SIRT1	SIRT3	Human Endogenous Metabolite
<b>In Vitro</b>	Nicotinamide riboside Chloride (0.5 nM; 24 hours) reduces the acetylation status of Ndufa9 and SOD2 <sup>[1]</sup> . Nicotinamide riboside Chloride increases intracellular and mitochondrial NAD <sup>+</sup> content in C2C12, Hepa1.6, and HEK293 cells in a concentration-dependent manner at concentrations ranging from 1-1000 μM <sup>[1]</sup> . Nicotinamide riboside chloride boosts NAD to restore antiviral poly(ADP-ribose) polymerase (PARP) functions to support innate immunity for coronavirus (CoVs), a cause of COVID-19 <sup>[3]</sup> .		

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Western Blot Analysis<sup>[1]</sup>

Cell Line:	HEK293T cells
Concentration:	0.5 nM
Incubation Time:	24 hours
Result:	Reduced the acetylation status of Ndufa9 and SOD2.

#### In Vivo

Chronic Nicotinamide riboside Chloride (p.o.; 400 mg/kg/day; for 16 weeks) supplementation increases plasma and intracellular NAD<sup>+</sup> content in a tissue-specific manner<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	10-week-old C57Bl/6J mice <sup>[1]</sup>
Dosage:	400 mg/kg
Administration:	PO; daily; for 16 weeks
Result:	Increased plasma and intracellular NAD <sup>+</sup> content in a tissue-specific manner.

## CUSTOMER VALIDATION

- Nat Commun. 2023 Jan 16;14(1):240.
- Mol Ther. 2022 Sep 21;S1525-0016(22)00567-6.
- Redox Biol. 2022 Oct 11;57:102507.
- Cell Biosci. 2021 Nov 10;11(1):192.
- Cells. 2023 Oct 2, 12(19), 2396.

See more customer validations on [www.MedChemExpress.com](http://www.MedChemExpress.com)

## REFERENCES

[1]. Cantó C, et al. The NAD(+) precursor nicotinamide riboside enhances oxidative metabolism and protects against high-fat diet-induced obesity. Cell Metab. 2012 Jun 6;15(6):838-47.

[2]. Bing Gong, et al. Nicotinamide Riboside Restores Cognition Through an Upregulation of Proliferator-Activated Receptor-γ Coactivator 1α Regulated β-Secretase 1 Degradation and Mitochondrial Gene Expression in Alzheimer's Mouse Models. Neurobiol Aging. 2013 Jun;34(6):1581-8.

[3]. Collin D Heer, et al. Coronavirus and PARP Expression Dysregulate the NAD Metabolome: A Potentially Actionable Component of Innate Immunity. bioRxiv. 2020 Apr 30;2020.04.17.047480.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA