

Product Data Sheet

Inhibitors

Screening Libraries

Proteins

OMDM-3

Molecular Weight:

Target: Endogenous Metabolite

Pathway: Metabolic Enzyme/Protease

453.66

Storage: Please store the product under the recommended conditions in the Certificate of

Analysis.

N OH OH

BIOLOGICAL ACTIVITY

Description	OMDM-3 is a selective and metabolically stable inhibitor of anandamide cellular uptake (ACU), with a K_i of 16.6 μ M $^{[1]}$.
IC ₅₀ & Target	Ki: 16.6 μ M (anandamide cellular uptake) $^{[1]}$
In Vitro	OMDM-3 shows poor affinity for either CB1 (K_i =6.1 μ M) or CB2 (K_i >10 μ M) receptors in rat brain and spleen membranes, respectively. OMDM-3 has almost no activity at vanilloid receptors in the intracellular calcium assay carried out with intact cells over-expressing the human VR1 (EC_{50} >10 μ M), and no activity as inhibitors of FAAH in N18TG2 cell membranes (K_i >50 μ M)[1]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Ortar G, et al. Novel selective and metabolically stable inhibitors of anandamide cellular uptake. Biochem Pharmacol. 2003 May 1;65(9):1473-81.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA