

## **Product** Data Sheet

## QD394

Cat. No.: HY-139369

CAS No.: 2132411-21-1

Molecular Formula:  $C_{19}H_{19}N_5O_2$ Molecular Weight: 349.39

Target: Reactive Oxygen Species

Pathway: Immunology/Inflammation; Metabolic Enzyme/Protease; NF-кВ

Storage: Please store the product under the recommended conditions in the Certificate of

Analysis.

## **BIOLOGICAL ACTIVITY**

| Description | QD394 is a reactive oxygen species (ROS) inducer that can induce lipid peroxidation, increase intracellular ROS accumulation, inhibit STAT3 phosphorylation, and induce ferroptosis <sup>[1]</sup> .   |
|-------------|--|
| In Vitro    | QD394 (0-10 $\mu$ M, 24 h) is cytotoxic with the IC $_{50}$ values of 0.64, 0.34 and 0.9 $\mu$ M for MIA PaCa-2, PANC-1 and BxPC-3 cell lines, respectively. And QD394 can inhibit the phosphorylation of STAT3 in cells and increase the phosphorylation of H2AX <sup>[1]</sup> . QD394(1-10 $\mu$ M, 4 h) mediates cell death by increasing cellular ROS levels and decreases GSH/GSSG ratio in MIA PaCa-2 cells; also induces iron-dependent and GPX4-mediated ferroptosis <sup>[1]</sup> . MCE has not independently confirmed the accuracy of these methods. They are for reference only. |

## **REFERENCES**

[1]. Shuai Hu, et al. A Novel Redox Modulator Induces a GPX4-Mediated Cell Death That Is Dependent on Iron and Reactive Oxygen Species. J Med Chem. 2020 Sep 10;63(17):9838-9855.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA