## (R)-ZG197

®

MedChemExpress

Cat. No.:	HY-152096	
Molecular Formula:	$C_{28}H_{35}F_{3}N_{4}O_{3}$	-
Molecular Weight:	532.6	
Target:	Bacterial; ClpP	F T H N
Pathway:	Anti-infection; Cell Cycle/DNA Damage	
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.	0,

BIOLOGICAL ACTIV			
Description	(R)-ZG197 is a highly selective Staphylococcus aureus Caseinolytic protease P ( <b>Sa</b> ClpP) activator with an EC <sub>50</sub> of 1.5 μM. (R)-ZG197 also activates Homo sapiens ClpP (HsClpP) with an EC <sub>50</sub> of 31.4 μM <sup>[1]</sup> .		
IC <sub>50</sub> & Target	EC50: 1.5 μM (SaClpP), 31.4 μM (HsClpP) <sup>[1]</sup> Kd: 2.5 μM (SaClpP) <sup>[1]</sup>		
In Vitro	<ul> <li>(R)-ZG197 (10 μM; 2 h) significantly enhances the thermal stability of SaClpP while having a weak effect on HsClpP<sup>[1]</sup>.</li> <li>(R)-ZG197 (0-256 μg/mL; 18 h) significantly suppresses S. aureus with a quantified MIC of 0.5 μg/mL. (R)-ZG197 displays strong antibacterial activity on a broad spectrum of S. aureus strains, with MIC values of 0.5-2 μg/mL<sup>[1]</sup>.</li> <li>(R)-ZG197 (0-20 μM) decrease SaFtsZ abundance in the 8325-4 S. aureus but not in the corresponding ΔclpP mutant strain<sup>[1]</sup> MCE has not independently confirmed the accuracy of these methods. They are for reference only.</li> <li>Western Blot Analysis<sup>[1]</sup></li> </ul>		
	Cell Line:	Cell lysates of S. aureus 8325-4 clpP knockout (ΔclpP) strain	
	Concentration:	0, 2.5, 5 and 10 μM	
	Incubation Time:	15 min	
	Result:	SaFtsZ protein was degraded when SaClpP was added.	
In Vivo	<ul> <li>(R)-ZG197 (25-100 mg/kg; i.p.; once) significantly prolong the survival rate in zebrafish USA300 infection model<sup>[1]</sup>.</li> <li>(R)-ZG197 (7.5 mg/kg; s.c.; twice a day for 3 days) shows anti-infective efficacy in murine skin S. aureus infection models<sup>[1]</sup>.</li> <li>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</li> </ul>		
	Animal Model:	Zebrafish USA300 infection model <sup>[1]</sup>	
	Dosage:	25, 50, or 100 mg/kg	
	Administration:	Intraperitoneal injection, single dose	
	Result:	Significantly prolong the survival rate at 50 mg/kg. Lost therapeutic effects on zebrafish infected with the $\Delta$ clpP mutant strain.	

Animal Model:	Female BALB/c mice, S. aureus infection model <sup>[1]</sup>
Dosage:	7.5 mg/kg
Administration:	Subcutaneous injection, twice a day for 3 days
Result:	Caused a smaller necrotic lesion size in mice compared with the vehicle control.

## REFERENCES

[1]. Wei B, et al. Anti-infective therapy using species-specific activators of Staphylococcus aureus ClpP. Nat Commun. 2022 Nov 14;13(1):6909.

## Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA