

Product Data Sheet

RXR antagonist 1

Cat. No.: HY-144377

Molecular Formula: $C_{28}H_{33}F_3N_2O_3$ Molecular Weight: 502.57

Target: RAR/RXR

Pathway: Metabolic Enzyme/Protease; Vitamin D Related/Nuclear Receptor

Storage: Powder -20°C 3 years

4°C 2 years

In solvent -80°C 6 months

-20°C 1 month

SOLVENT & SOLUBILITY

In Vitro

DMSO: 100 mg/mL (198.98 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	1.9898 mL	9.9489 mL	19.8977 mL
	5 mM	0.3980 mL	1.9898 mL	3.9795 mL
	10 mM	0.1990 mL	0.9949 mL	1.9898 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: 2.5 mg/mL (4.97 mM); Clear solution; Need ultrasonic
- 2. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: 2.5 mg/mL (4.97 mM); Clear solution; Need ultrasonic

BIOLOGICAL ACTIVITY

Pescription RXR antagonist 1 (compound 6a) is a retinoid X receptor (RXR) modulator. RXR antagonist 1 shows potent RXR-antagonistic

activity, with a pA₂ of 8.06. RXR antagonist 1 can be used for type 2 diabetes research^[1].

In Vitro RXR antagonist 1 (compound 6a) shows potent RXR-antagonistic activities at $1 \mu M^{[1]}$.

RXR antagonist 1 shows competitive binding to the LBP in hRXR α -LBD, with K_i of 0.384 \pm 0.072, K_d of 0.277 \pm 0.038, and K_i/K_d

of 1.39^[1].

 $\label{thm:continuous} The cell permeability of RXR antagonist 1 shows no correlation with RXR-antagonistic activity \emph{$[1]$}.$

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES



Page 2 of 2 www.MedChemExpress.com