

Product Data Sheet

Reveromycin A

Cat. No.: HY-129337 CAS No.: 134615-37-5 Molecular Formula: $C_{36}H_{52}O_{11}$ Molecular Weight: 660.79

Target:Antibiotic; Apoptosis; FungalPathway:Anti-infection; ApoptosisStorage:Powder -20°C 3 years

In solvent

-80°C 6 months -20°C 1 month

poptosis; Fungal n; Apoptosis -20°C 3 years

BIOLOGICAL ACTIVITY

Description	Reveromycin A, a benzoquinoid antibiotic isolated from the genus Streptomyces, is a selective inhibitor of protein synthesis in eukaryotic cells. Reveromycin A inhibits bone resorption by inducing apoptosis specifically in osteoclasts. Reveromycin A has antiproliferative activity against tumor cell lines and antifungal activity ^{[1][2]} .
In Vitro	Reveromycin A inhibits protein synthesis in osteoclasts (OCs) by selectively blocking enzymatic activity of isoleucyl-tRNA synthetase ^[2] . Reveromycin A (0.3-30 μ M; for 24 h) induces apoptosis in OCs through inhibition of protein synthesis ^[2] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.
In Vivo	Reveromycin A (20 mg/kg; twice daily; i.v.; for 3 days) significantly decreases numbers of OCs without affecting numbers of osteoblasts in normal 4-week-old male Sprague-Dawley rats ^[2] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

 $[1]. \ H\ Takahashi\ , et\ al.\ Reveromycins, new\ inhibitors\ of\ eukaryotic\ cell\ growth.\ II.\ Biological\ activities.\ J\ Antibiot\ (Tokyo).\ 1992\ Sep; 45(9):1414-9.$

[2]. Je-Tae Woo, et al. Reveromycin A, an agent for osteoporosis, inhibits bone resorption by inducing apoptosis specifically in osteoclasts. Proc Natl Acad Sci U S A. 2006 Mar 21;103(12):4729-34.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA