## Sonlicromanol hydrochloride

 Cat. No.:
 HY-120332

 CAS No.:
 2162149-24-6

 Molecular Formula:
  $C_{19}H_{29}ClN_2O_3$ 

Molecular Weight: 368.9

Target: Reactive Oxygen Species

Pathway: Immunology/Inflammation; Metabolic Enzyme/Protease; NF-κΒ

**Storage:** 4°C, protect from light, stored under nitrogen

\* In solvent: -80°C, 6 months; -20°C, 1 month (protect from light, stored under

nitrogen)

## **SOLVENT & SOLUBILITY**

In Vitro DMSO: 170 mg/mL (460.83 mM; Need ultrasonic)

 $H_2O : \ge 100 \text{ mg/mL} (271.08 \text{ mM})$ 

\* "≥" means soluble, but saturation unknown.

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	2.7108 mL	13.5538 mL	27.1076 mL
	5 mM	0.5422 mL	2.7108 mL	5.4215 mL
	10 mM	0.2711 mL	1.3554 mL	2.7108 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo 1. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 5 mg/mL (13.55 mM); Clear solution

2. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline

Solubility: ≥ 4.25 mg/mL (11.52 mM); Clear solution

3. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline)

Solubility: ≥ 4.25 mg/mL (11.52 mM); Clear solution

## **BIOLOGICAL ACTIVITY**

Description Sonlicromanol (KH176) hydrochloride, a chemical entity derivative of Trolox, is a blood-brain barrier permeable ROS-redox modulator. Sonlicromanol (KH176) hydrochloride is used in the study for mitochondrial disorders<sup>[1]</sup>.

In Vivo Sonlicromanol (KH176) hydrochloride maintains microstructural coherence in the brain of Ndufs4<sup>-/-</sup> mice<sup>[1]</sup>.

 $\label{eq:mce} \mbox{MCE has not independently confirmed the accuracy of these methods. They are for reference only.}$ 

Animal Model:	Ndufs4 $^{-/-}$ mice (Leigh Disease model) $^{[1]}$ .		
Dosage:	10 mg/kg.		
Administration:	IP, daily (PD14-PD45, 32 days).		
Result:	Significantly improve rotarod and gait performance and reduced the degeneration of retinal ganglion cells.  Resulted in statistically significantly higher FA values in the external capsule and a similar trend was found in the cerebral peduncle.		

## **REFERENCES**

[1]. Ria de Haas, et al. Therapeutic effects of the mitochondrial ROS-redox modulator KH176 in a mammalian model of Leigh Disease. Sci Rep. 2017 Sep 15;7(1):11733.

 $\label{lem:caution:Product} \textbf{Caution: Product has not been fully validated for medical applications. For research use only.}$ 

Tel: 609-228-6898

Fax: 609-228-5909

 $\hbox{E-mail: tech@MedChemExpress.com}$ 

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA