


TAK-960 dihydrochloride

Cat. No.:	HY-15160B
Molecular Formula:	C ₂₇ H ₃₆ Cl ₂ F ₃ N ₇ O ₃
Molecular Weight:	634.52
Target:	Polo-like Kinase (PLK)
Pathway:	Cell Cycle/DNA Damage
Storage:	4°C, sealed storage, away from moisture * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)



SOLVENT & SOLUBILITY

In Vitro

H₂O : 10 mg/mL (15.76 mM; Need ultrasonic)
DMSO : 3.23 mg/mL (5.09 mM; Need ultrasonic)

Solvent	Mass	Concentration		
		1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	1.5760 mL	7.8800 mL	15.7599 mL
	5 mM	0.3152 mL	1.5760 mL	3.1520 mL
	10 mM	0.1576 mL	0.7880 mL	1.5760 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description

TAK-960 dihydrochloride is an orally available, selective inhibitor of polo-like kinase 1 (PLK1), with an IC₅₀ of 0.8 nM. TAK-960 dihydrochloride also shows inhibitory activities against PLK2 and PLK3, with IC₅₀s of 16.9 and 50.2 nM, respectively. TAK-960 dihydrochloride inhibits proliferation of multiple cancer cell lines and exhibits significant efficacy against multiple tumor xenografts^[1].

IC₅₀ & Target

PLK1 0.8 nM (IC ₅₀)	PLK2 16.9 nM (IC ₅₀)	PLK3 50.2 nM (IC ₅₀)	FAK/PTK2 19.6 nM (IC ₅₀)
MLCK/MYLK 25.6 nM (IC ₅₀)	FES/FPS 58.2 nM (IC ₅₀)		

In Vitro

TAK-960 dihydrochloride treatment causes accumulation of G2-M cells, aberrant polo mitosis morphology, and increased phosphorylation of histone H3 (pHH3). TAK-960 dihydrochloride (2-1000 nM; 72 hours) inhibits proliferation of multiple cancer cell lines, with mean EC₅₀ values ranging from 8.4 to 46.9 nM, but not in nondividing normal cells^[1]. TAK-960 dihydrochloride (8 nM) leads to G2/M cell cycle arrest without significant cytotoxicity in HeLa cells^[2]. MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Cell Viability Assay^[1]

Cell Line:	HT-29, HCT116, COLO320DM, HCT-15, RKO, SW480, K-562...Hela, DU 145 cells
Concentration:	2-1000 nM
Incubation Time:	72 hours
Result:	Inhibited proliferation of human cancer cell lines regardless of TP53 and KRAS mutation and MDR1 expression status.

In Vivo

TAK-960 dihydrochloride exhibits (10 mg/kg; p.o.; once daily for 2 weeks) significant efficacy against multiple tumor xenografts^[1].
In animal models, TAK-960 dihydrochloride (p.o.) increases pHH3 in a dose-dependent manner and significantly inhibits the growth of HT-29 colorectal cancer xenografts^[1].
MCE has not independently confirmed the accuracy of these methods. They are for reference only.

Animal Model:	nude mice or SCID mice (bearing HCT116, PC-3, BT474, A549, NCI-H1299, NCI-H1975, A2780, and MV4-11 cells) ^[1]
Dosage:	10 mg/kg
Administration:	P.o.; once daily for 2 weeks
Result:	Substantial antitumor activity and good tolerability.

CUSTOMER VALIDATION

- Sci Transl Med. 2018 Jul 18;10(450):eaaq1093.
- Cancer Lett. 2020 Oct 28;491:50-59.

See more customer validations on www.MedChemExpress.com

REFERENCES

[1]. Hikichi Y, et al. TAK-960, a novel, orally available, selective inhibitor of polo-like kinase 1, shows broad-spectrum preclinical antitumor activity in multiple dosing regimens. Mol Cancer Ther. 2012 Mar;11(3):700-9.

[2]. Inoue M, et al. PLK1 blockade enhances therapeutic effects of radiation by inducing cell cycle arrest at the mitotic phase. Sci Rep. 2015 Oct 27;5:15666.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA