TD-165

Cat. No.:	HY-130714
CAS No.:	2305936-56-3
Molecular Formula:	C ₄₆ H ₅₉ N ₇ O ₈ S
Molecular Weight:	870.07
Target:	PROTACs
Pathway:	PROTAC
Storage:	-20°C, protect from light
	* In solvent : -80°C, 6 months; -20°C, 1 month (protect from light)

SOLVENT & SOLUBILITY

		Solvent Mass Concentration	1 mg	5 mg	10 mg
Preparing Stock Solutions	Preparing Stock Solutions	1 mM	1.1493 mL	5.7467 mL	11.4933 mL
		5 mM	0.2299 mL	1.1493 mL	2.2987 mL
		10 mM	0.1149 mL	0.5747 mL	1.1493 mL

BIOLOGICAL ACTI	ЛТҮ	
Description		ed cereblon (CRBN) degrader. TD-165 comprises a cereblon (CRBN) ligand binding group, a linker au (VHL) binding group ^[1] .
IC₅₀ & Target	Cereblon 20.4 nM (DC50)	
In Vitro	nM and 99.6%, respectiv TD-165 (0.1-10 μM; 24 hc	urs) shows 50% degradation concentration (DC ₅₀) and maximum degradation (D _{max}) value of 20.4 rely ^[1] . Dours) leads to a concentration-dependent decrease in the level of CRBN proteins in HEK293T cells ^[1] . ntly confirmed the accuracy of these methods. They are for reference only.
	Cell Line:	HEK293T cells
	Concentration:	0-100 nM
	Incubation Time:	24 hours

Product Data Sheet

Result:	Exhibited an DC ₅₀ value of 20.4 nM.
Western Blot Analysis ^[1]	
Cell Line:	HEK293T cells
Concentration:	0.1, 1, and 10 μM
Incubation Time:	24 hours
Result:	Decreased the level of CRBN protein.

REFERENCES

[1]. Kim K, et al. Disordered region of cereblon is required for efficient degradation by proteolysis-targeting chimera. Sci Rep. 2019 Dec 23;9(1):19654.

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA