


## TH1834

<b>Cat. No.:</b>	HY-123604
<b>CAS No.:</b>	2108830-08-4
<b>Molecular Formula:</b>	C <sub>33</sub> H <sub>40</sub> N <sub>6</sub> O <sub>3</sub>
<b>Molecular Weight:</b>	568.71
<b>Target:</b>	Histone Acetyltransferase; Apoptosis
<b>Pathway:</b>	Epigenetics; Apoptosis
<b>Storage:</b>	Powder    -20°C    3 years 4°C        2 years In solvent   -80°C    6 months -20°C    1 month



### SOLVENT & SOLUBILITY

<b>In Vitro</b>	DMSO : 10 mg/mL (17.58 mM; ultrasonic and warming and heat to 60°C)				
		Solvent Concentration	Mass 1 mg	5 mg	10 mg
	<b>Preparing Stock Solutions</b>	1 mM	1.7584 mL	8.7918 mL	17.5837 mL
		5 mM	0.3517 mL	1.7584 mL	3.5167 mL
10 mM		0.1758 mL	0.8792 mL	1.7584 mL	
Please refer to the solubility information to select the appropriate solvent.					
<b>In Vivo</b>	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 1 mg/mL (1.76 mM); Clear solution 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 1 mg/mL (1.76 mM); Clear solution 3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 1 mg/mL (1.76 mM); Clear solution				

### BIOLOGICAL ACTIVITY

<b>Description</b>	TH1834 is a specific Tip60 (KAT5) histone acetyltransferase (HAT) inhibitor. TH1834 induces apoptosis and increases DNA damage in breast cancer. TH1834 does not affect the activity of related histone acetyltransferase MOF. Anticancer activity <sup>[1]</sup> .
<b>IC<sub>50</sub> &amp; Target</b>	TIP60
<b>In Vitro</b>	TH1834 (0-500 μM; 1 hour; MCF7 cells) treatment significantly reduces the viability of MCF7 cells <sup>[1]</sup> . TH1834 (0-500 μM; 1 hour; MCF7 cells) treatment significantly increases cytotoxicity in MCF7 cells <sup>[1]</sup> .

TH1834 (500  $\mu$ M; 1 hour; MCF7 cells) treatment induces caspase 3 activation in MCF7 cells<sup>[1]</sup>.

TH1834 significantly inhibits Tip60 activity in vitro and treating cells with TH1834 results in apoptosis and increased unrepaired DNA damage in breast cancer<sup>[1]</sup>.

MCE has not independently confirmed the accuracy of these methods. They are for reference only.

#### Cell Viability Assay<sup>[1]</sup>

Cell Line:	MCF7 cells
Concentration:	0 $\mu$ M, 0.5 $\mu$ M, 5 $\mu$ M, 50 $\mu$ M and 500 $\mu$ M
Incubation Time:	1 hour
Result:	Significantly reduced the viability of MCF7 cells.

#### Cell Cytotoxicity Assay<sup>[1]</sup>

Cell Line:	MCF7 cells
Concentration:	0 $\mu$ M, 0.5 $\mu$ M, 5 $\mu$ M, 50 $\mu$ M and 500 $\mu$ M
Incubation Time:	1 hour
Result:	Highly significant increase in cytotoxicity at all concentrations used.

#### Western Blot Analysis<sup>[1]</sup>

Cell Line:	MCF7 cells
Concentration:	500 $\mu$ M
Incubation Time:	1 hour
Result:	Marked caspase 3 activation was observed in MCF7 cells in an independent assay.

## REFERENCES

[1]. Gao C, et al. Rational design and validation of a Tip60 histone acetyltransferase inhibitor. Sci Rep. 2014 Jun 20;4:5372.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA