Product Data Sheet

Tos-PEG2-NH-Boc

Cat. No.: HY-135798 CAS No.: 192132-77-7 Molecular Formula: $C_{16}H_{25}NO_{6}S$ Molecular Weight: 359.44

Target: **PROTAC Linkers**

Pathway: **PROTAC**

Storage: Powder -20°C 3 years

 $4^{\circ}C$ 2 years

In solvent -80°C 6 months

> -20°C 1 month

SOLVENT & SOLUBILITY

In Vitro

DMSO: 250 mg/mL (695.53 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	2.7821 mL	13.9105 mL	27.8211 mL
	5 mM	0.5564 mL	2.7821 mL	5.5642 mL
	10 mM	0.2782 mL	1.3911 mL	2.7821 mL

Please refer to the solubility information to select the appropriate solvent.

In Vivo

- 1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.08 mg/mL (5.79 mM); Clear solution
- 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (5.79 mM); Clear solution
- 3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.08 mg/mL (5.79 mM); Clear solution

BIOLOGICAL ACTIVITY

Description	${\it Tos-PEG2-NH-Boc}\ is\ a\ {\it PEG-based}\ {\it PROTAC}\ linker\ that\ can\ be\ used\ in\ the\ synthesis\ of\ {\it PROTACs}^{[1]}.$		
IC ₅₀ & Target	PEGs	Alkyl/ether	
In Vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.		

FERENCES	
An S, et al. Small-molec	cule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562.
	Caution: Product has not been fully validated for medical applications. For research use only.
	Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com
	Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 2 of 2 www.MedChemExpress.com