


Tubulin inhibitor 7

Cat. No.:	HY-136122
CAS No.:	1309925-41-4
Molecular Formula:	C ₂₁ H ₁₄ N ₂ O ₄
Molecular Weight:	358.35
Target:	Microtubule/Tubulin
Pathway:	Cell Cycle/DNA Damage; Cytoskeleton
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Tubulin inhibitor 7 (Compound 33c) is a tubulin inhibitor and a potent inhibitor of multiple cancer cell lines. Tubulin inhibitor 7 inhibits tubulin polymerization with an IC ₅₀ of 0.52 μM. Tubulin inhibitor 7 inhibits K562 cell growth with an IC ₅₀ of 11 nM ^[1] .
IC₅₀ & Target	IC ₅₀ : 0.52 μM (tubulin polymerization) ^[1]
In Vitro	Tubulin inhibitor 7 blocks mitosis through an arrest of cells in the G ₂ /M phase, as illustrated in typical histograms ^[1] . Tubulin inhibitor 7 shows excellent antiproliferative potencies with IC ₅₀ s of 40, 24, 16, 6, 29, and 16 nM for NCIH460, SKOV3, BT549, 451LU, SW480, and DLD-1 tumor cell lines, respectively ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Prinz H, et al. N-benzoylated phenoxazines and phenothiazines: synthesis, antiproliferative activity, and inhibition of tubulin polymerization. J Med Chem. 2011 Jun 23;54(12):4247-63.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA