


Tubulin inhibitor 8

Cat. No.:	HY-136123
CAS No.:	1309925-39-0
Molecular Formula:	C ₂₁ H ₁₄ N ₂ O ₃
Molecular Weight:	342.35
Target:	Microtubule/Tubulin
Pathway:	Cell Cycle/DNA Damage; Cytoskeleton
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Tubulin inhibitor 8 (Compound 33b) is a tubulin inhibitor and a potent inhibitor of multiple cancer cell lines. Tubulin inhibitor 8 inhibits tubulin polymerization with an IC ₅₀ of 0.73 μM. Tubulin inhibitor 8 inhibits K562 cell growth with an IC ₅₀ of 14 nM ^[1] .
IC₅₀ & Target	IC ₅₀ : 0.73 μM (tubulin polymerization) ^[1]
In Vitro	Tubulin inhibitor 8 blocks mitosis through an arrest of cells in the G ₂ /M phase, as illustrated in typical histograms ^[1] . Tubulin inhibitor 8 shows excellent antiproliferative potencies with IC ₅₀ s of 15, 6, 8, 2, 8, 6, and 9 nM for NCIH460, SKOV3, BT549, 451LU, SW480, COLO-205, and DLD-1 tumor cell lines, respectively ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

[1]. Prinz H, et al. N-benzoylated phenoxazines and phenothiazines: synthesis, antiproliferative activity, and inhibition of tubulin polymerization. J Med Chem. 2011 Jun 23;54(12):4247-63.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA