(+)-Biotin-PEG6-hydrazide

MedChemExpress

Cat. No.:	HY-130891	
Molecular Formula:	C ₂₅ H ₄₇ N ₅ O ₉ S	
Molecular Weight:	593.73	
Target:	PROTAC Linkers	$\overset{O_{\frac{1}{2}} \to WH}{\underset{H}{\overset{WL}{}}} \overset{O_{\frac{1}{2}}}{\overset{WH}{}} \overset{O_{\frac{1}{2}}}{\overset{O_{\frac{1}{2}}}} \overset{O_{\frac{1}{2}}}{} \overset{O_{\frac{1}{2}}}{} \overset{O_{\frac{1}{2}}}{} \overset{O_{\frac{1}{2}}} \overset{O_{\frac{1}{2}}}{} \overset{O_{\frac{1}{2}}}} \overset{O_{\frac{1}{2}}}{} \overset{O_{\frac{1}{2}}}}{} \overset{O_{\frac{1}{2}}}} $
Pathway:	PROTAC	
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.	

BIOLOGICAL ACTIVITY		
Description	(+)-Biotin-PEG6-hydrazide is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs ^[1] .	
IC ₅₀ & Target	PEGs	
In Vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.	

REFERENCES

[1]. An S, et al. Small-molecule PROTACs: An emerging and promising approach for the development of targeted therapy drugs. EBioMedicine. 2018 Oct;36:553-562.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909 E-mail: tech@MedChemExpress.com Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Product Data Sheet