


Sacubitril-¹³C₄ hemicalcium salt

Cat. No.:	HY-15407AS
Molecular Formula:	C ₂₀ ¹³ C ₄ H ₂₈ Ca _{0.5} NO ₅
Molecular Weight:	434.49
Target:	Nepriylsin; Isotope-Labeled Compounds
Pathway:	Metabolic Enzyme/Protease; Others
Storage:	Please store the product under the recommended conditions in the Certificate of Analysis.



BIOLOGICAL ACTIVITY

Description	Sacubitril- ¹³ C ₄ (hemicalcium salt) is a ¹³ C-labeled and deuterium labeled Sacubitril hemicalcium salt. Sacubitril (AHU-377) hemicalcium salt is a potent NEP inhibitor with an IC ₅₀ of 5 nM. Sacubitril hemicalcium salt is a component of the heart failure medicine LCZ696[1].
In Vitro	Stable heavy isotopes of hydrogen, carbon, and other elements have been incorporated into drug molecules, largely as tracers for quantitation during the drug development process. Deuteration has gained attention because of its potential to affect the pharmacokinetic and metabolic profiles of drugs ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Russak EM, et al. Impact of Deuterium Substitution on the Pharmacokinetics of Pharmaceuticals. *Ann Pharmacother.* 2019;53(2):211-216.
- [2]. Ksander GM, et al. Dicarboxylic acid dipeptide neutral endopeptidase inhibitors. *J Med Chem.* 1995 May 12;38(10):1689-700.;Voors AA, et al. The potential role of valsartan + AHU377 (LCZ696) in the treatment of heart failure. *Expert Opin Investig Drugs.* 2013 Aug;22(8):1041-7.;von Lueder TG, et al. Angiotensin receptor neprilysin inhibitor LCZ696 attenuates cardiac remodeling and dysfunction after myocardial infarction by reducing cardiac fibrosis and hypertrophy. *Circ Heart Fail.* 2015 Jan;8(1):71-8.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA