D-3263 hydrochloride

Cat. No.:	HY-16162A	\frown
CAS No.:	1008763-54-9	
Molecular Formula:	C ₂₁ H ₃₂ ClN ₃ O ₃	·**=0
Molecular Weight:	409.95	Ň
Target:	TRP Channel	
Pathway:	Membrane Transporter/Ion Channel; Neuronal Signaling	
Storage:	4°C, sealed storage, away from moisture * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)	NH₂ H−Cl

SOLVENT & SOLUBILITY

		Solvent Mass Concentration	1 mg	5 mg	10 mg		
	Preparing Stock Solutions	1 mM	2.4393 mL	12.1966 mL	24.3932 mL		
		5 mM	0.4879 mL	2.4393 mL	4.8786 mL		
		10 mM	0.2439 mL	1.2197 mL	2.4393 mL		
	Please refer to the so	olubility information to select the ap	propriate solvent.	1			
In Vivo		1. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 14 mg/mL (34.15 mM); Clear solution					
		2. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 3.5 mg/mL (8.54 mM); Clear solution					
		3. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 3.5 mg/mL (8.54 mM); Clear solution					

BIOLOGICAL ACTIVITY				
In Vitro	D-3263 hydrochloride binds to and activates TRPM8, which may result in an increase in calcium and sodium entry; the disruption of calcium and sodium homeostasis; and the induction of cell death in TRPM8-expressing tumor cells. D-3263 hydrochloride may decrease dihydrotestosterone (DHT) levels, which may contribute to its inhibitory effects on prostate cancer and BPH ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.			



REFERENCES

[1]. enteric-coated TRPM8 agonist D-3263 hydrochloride

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA