HDAC-IN-3

Cat. No.:	HY-19772				
CAS No.:	1018673-42-	1			
Molecular Formula:	C ₂₂ H ₃₃ N ₃ O ₄				
Molecular Weight:	403.52				
Target:	HDAC				
Pathway:	Cell Cycle/DNA Damage; Epigenetics				
Storage:	Powder	-20°C	3 years		
		4°C	2 years		
	In solvent	-80°C	2 years		
		-20°C	1 year		

SOLVENT & SOLUBILITY

In Vitro	DMSO : 50 mg/mL (123.91 mM; Need ultrasonic)						
	Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg		
		1 mM	2.4782 mL	12.3910 mL	24.7819 mL		
		5 mM	0.4956 mL	2.4782 mL	4.9564 mL		
		10 mM	0.2478 mL	1.2391 mL	2.4782 mL		
	Please refer to the solubility information to select the appropriate solvent.						
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: ≥ 2.5 mg/mL (6.20 mM); Clear solution						
	2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.5 mg/mL (6.20 mM); Clear solution						
	3. Add each solvent one by one: 10% DMSO >> 90% corn oil Solubility: ≥ 2.5 mg/mL (6.20 mM); Clear solution						

DLOGICAL ACTIV	ТҮ
Description	HDAC-IN-3 is a histone deacetylase (HDAC) inhibitor, extracted from patent WO/2008040934 A1.Target: HDAC
IC ₅₀ & Target	HDAC

REFERENCES

Product Data Sheet





[1]. Davidson Alan Hornsby, et al. Preparation of amino acid-based hydroxamic acids as histone deacetylase (HDAC) inhibitors. From PCT Int. Appl. (2008), WO 2008040934 A1 20080410.

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA