Mal-PEG-OH (MW 5000)

Target:BiochemPathway:Others	H-9 hC ₉ H ₁₂ N ₂ O ₄ hical Assay Reagents tore the product under the recommended conditions in the Certificate of
------------------------------	---

Description Mal-PEG-OH (MW 5000) can be used as a macroinitiator to obtain amphiphilic diblock copolymers by ring-opening polymerization of LA. Nanoparticles prepared using amphiphilic block copolymers can form active drug delivery systems. Nanoparticles encapsulate Triptolide (HY-32735), which can avoid the disadvantage of Triptolide's poor water solubility and	BIOLOGICAL ACTIVITY	
reduce its toxicity.	Description	polymerization of LA. Nanoparticles prepared using amphiphilic block copolymers can form active drug delivery systems. Nanoparticles encapsulate Triptolide (HY-32735), which can avoid the disadvantage of Triptolide's poor water solubility and

REFERENCES

[1]. Chen XY, et al. The potential of follicle-stimulating hormone peptide-modified triptolide-loaded nanoparticles to induce a mouse model of premature ovarian insufficiency. Int J Nanomedicine. 2015 Apr 7;10:2765-74.

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA



Product Data Sheet