


Niperotidine

Cat. No.:	HY-15539
CAS No.:	84845-75-0
Molecular Formula:	C ₂₀ H ₂₆ N ₄ O ₅ S
Molecular Weight:	434.51
Target:	Histamine Receptor
Pathway:	GPCR/G Protein; Immunology/Inflammation; Neuronal Signaling
Storage:	4°C, sealed storage, away from moisture and light * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light)



SOLVENT & SOLUBILITY

In Vitro	DMSO : 100 mg/mL (230.14 mM; Need ultrasonic)				
		Solvent Concentration	Mass		
	Preparing Stock Solutions		1 mg	5 mg	10 mg
		1 mM	2.3014 mL	11.5072 mL	23.0144 mL
		5 mM	0.4603 mL	2.3014 mL	4.6029 mL
	10 mM	0.2301 mL	1.1507 mL	2.3014 mL	
Please refer to the solubility information to select the appropriate solvent.					
In Vivo	1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline Solubility: 2.5 mg/mL (5.75 mM); Suspended solution; Need ultrasonic 2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE-β-CD in saline) Solubility: 2.5 mg/mL (5.75 mM); Suspended solution; Need ultrasonic				

BIOLOGICAL ACTIVITY

Description	Niperotidine is a histamine H2-receptor antagonist.
IC₅₀ & Target	H ₂ Receptor
In Vivo	Niperotidine (piperonyl-ranitidine) is a H2 blocking agent. Niperotidine is a H2-receptor antagonist structurally related to ranitidine. After oral administration, it reaches a plasmatic peak within 60-120 min and is eliminated either in the urine or in the faeces, with an enterohepatic circulation ^[2] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

- [1]. Palasciano G, et al. The effect of the H₂-antagonist Niperotidine on intragastric acidity in healthy subjects undergoing 24-hour pH-monitoring. *Ital J Gastroenterol.* 1990 Oct;22(5):291-4.
- [2]. Gasbarrini G, et al. Acute liver injury related to the use of Niperotidine. *J Hepatol.* 1997 Sep;27(3):583-6.
-

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA